


STUDENT'S

Pocket Prescriber



22102052899

Med

K14094

mf

1/10/20

THE STUDENTS'
POCKET PRESCRIBER
AND
GUIDE TO PRESCRIPTION WRITING

BY
H. AUBREY HUSBAND, M.B.,
C.M., B.Sc., F.R.C.S.E., M.R.C.S.

FOURTH EDITION
Revised and Enlarged

EDINBURGH:
E. & S. LIVINGSTONE

1914

WELLCOME INSTITUTE
LIBRARY

Coll.

welMomec


Call

No.

151

CONTENTS

	Page
Introduction	5
Prescriptions for Alimentary Diseases	11
„ Blood Diseases	29
„ Circulatory Diseases	31
„ Constitutional Diseases	36
„ Ear, Eye, Nose, and Throat Diseases	41
„ Infectious Diseases	45
„ Intoxications	57
„ Nervous Diseases	60
„ Respiratory Diseases	68
„ Skin Diseases	85
„ Urinary Diseases	101
„ Uterine and Genital Diseases	106
„ Children's Diseases	112
Appendix :—	
Tables of Weights and Measures—	
Imperial and Metric	128
Tables of Conversion of Imperial into Metric, and <i>vice versa</i>	129
Table of Strength of Solution	131
Incompatibilities	131.
Numerals (Latin)	133
Quantities (Latin)	135
Vocabulary of Latin Words and Phrases	137
Abbreviations	146
Index	148



INTRODUCTION.

EXAMINERS now generally require the student to write his prescriptions in unabbreviated Latin for medical qualifications, and for this reason most of the formulæ in this little book have been fully printed in that language. Even if he discard the usage in after days, it is necessary that he should be acquainted with the recognised forms in which prescriptions are set forth.

Most of the prescriptions which appeared in former editions have been retained, but a number of new ones have been introduced, derived from reliable sources, and embracing a more modern type of therapeutic remedies. It will be noted, however, that few of the drugs so largely advertised are included, many of which owe their popularity to successful puffing on the part of manufacturers; and the student must be on his guard against the specious and sometimes unscrupulous methods of drug agencies in palming off remedies which are often of no therapeutic value.

A medical prescription has a more or less recognised form. It consists of four parts—1. Superscription; 2. Inscription; 3. Subscription; and 4. Signature.

1. **Superscription.**—This is merely the Latin word "Recipe" contracted to the initial letter, with a stroke passed through the descending line thus, R, and means "Take thou."

2. **Inscription.**—Consisting of the drug or drugs used in compounding the mixture or other preparation. The quantities are written in the accusative case, and the names in the genitive, with the exception of the vehicle in which these are to be dissolved, suspended, or mixed, which may be either in the accusative or genitive, opinions being divided as to the most suitable case.

3. **Subscription.**—This is preceded by the word "Signa" or "Signetur," and constitutes the directions to the patient as to the amount, mode, and time of taking or using the preparation. Though written in Latin, it is translated into English on the label by the dispenser. It is an essential part of the prescription, and should never be omitted.

4. **Signature.**—This consists usually of the initials of the physician, but in some cases the full signature is appended. The latter practice is to be recommended, as it renders it more difficult to copy and use the prescription for others than the patient for whom it was originally given—a dangerous and reprehensible practice which is frequently adopted.

It has been thought desirable in this edition to alter the arrangement of the prescriptions by grouping them according to the disease for which they have been prescribed, in place of the actions which they are supposed to possess, and also to classify them under the system to which the diseases belong. This has been found more serviceable by those who have made use of such books, since drugs have often more than one action, and in the case of prescriptions which contain several drugs, may have opposite actions. It has also been thought advisable to group the prescriptions for children separately.

The quantities of the drugs used are expressed by the symbols and numerals used in both the Imperial and the Metric Standards of weights and measures, the latter being approximate only. It is hoped that this will make the book more useful to the student who intends to practise abroad.

In writing a prescription there are four simple rules which should be carried in the memory. In compounding medicines, according to Dr Paris in his *Pharmacologia*, we seek

1. **To Augment, Correct, or Modify the Action of a Medicine.**—Thus a combination of

purgatives is advisable especially if they have each an action on different parts of the alimentary canal. Ipecacuanha increases the purgative action of jalap and the emetic action of tartar emetic. The griping tendency of some drugs is modified by aromatics, and mercury is prevented from running off by the bowels by combining it with opium.

2. **To obtain the Joint Operation of two or more Medicines.**—The combination of purgatives just instanced will serve as an example of this rule, and also the combination of vaso-dilators and cardiac tonics in treating dilatation of the heart.

3. **To obtain a new Remedy.**—A combination of the tinctura ferri perchlor. and solution ammon. acet., of ferri sulph. with potass carb., and of acids and alkalis in effervescent preparations are familiar examples.

4. **To afford a Suitable Form.**—Glycerine of tracacanth and extracts for pills, emulsion of acacia or tragacanth or syrups to suspend heavy powders, etc. Glycerine disguises the taste of the tincture of iron; the essential oil of lemons that of the sulphate of magnesia.

The prescriber must be on his guard against the over-use of combinations, as in the blunderbuss prescriptions of the past, which aimed at a multitude of objects with the same charge, in the hope that one might perchance hit the mark. It should not be forgotten that treatment is more and more aiming at the removal of the cause of disease in place of relief of symptoms. Prophylaxis is taking the place of therapy, and empiricism is regarded as the refuge of ignorance. Care should be taken to make medicine as palatable as possible, especially for children.

In prescribing medicines for the sick, several important points have to be considered. The age, sex, race, temperament, and idiosyncrasy of the patient, the climate and disease, the mode and time of administration, and the compatibilities of the

drugs used in compounding the medicine, must all claim attention.

Let us briefly take in turn some of the points mentioned.

First, as to **Age**. This naturally influences the dose, and for our guidance several rules have been laid down. For instance, it has been proposed to add twelve to the age, and then to divide the age by the result. A child one year old would thus require a dose the thirteenth of that required for an adult; a child three years old, three-fifteenths, or one fifth. Another rule, equally useful, is to take the age at the next birthday and divide it by twenty-four; thus the dose for one year is found by taking the age of the child at two years and dividing it by twenty-four; for example, one year $\frac{2}{24} = \frac{1}{12}$, two years $\frac{3}{24} = \frac{1}{8}$, three years $\frac{4}{24} = \frac{1}{6}$, &c. We must also not forget that children bear narcotics badly, and that, as a rule, opium should be avoided in very young children, although, on the other hand, they tolerate belladonna and hyoscyamus very well. The rule for the administration of opium is one drop of the tincture for each year of age.

With regard to **Sex**, women, as a rule, require smaller doses than men, and this is especially the case with regard to purgatives; powerful cathartics, from the straining they cause, may give rise to displacement of the pelvic organs, and during pregnancy, to abortion.

The influence of race in respect to the actions of drugs has not been studied systematically, but it is believed that considerable differences exist between people of alien race in this respect, as well as in their susceptibility to, and the severity of attack in certain diseases.

The **Temperament** and **Idiosyncrasy** of individuals frequently seem to be outside the reign of all law, and can only be known by experiment. but the effects of habit are better understood. Indulgence in alcohol, opium, and in arsenic eating.

renders the system of those who thus indulge so tolerant of their action, that doses poisonous when administered to others have little or no effect on them.

The effect of climate, especially with regard to the action of narcotic and anti-periodic remedies, has been noted in the Tropics as compared with the temperate zone, and high atmospheric temperature probably increases the energy of many drugs.

The effect of **Disease** may be seen in the profuse salivation produced by even small doses of mercury in Bright's disease of the kidney, and the diminished influence of narcotics in delirium tremens, and in severe pain from cancer, the passage of gall stones, etc. The risk incurred by the administration of opium in erysipelas of the head, or congestion of the brain, or of purgatives in intussusception of the bowels, are all cases in point.

The question of the **Mode** and **Time** of administration of medicines demands more attention than is generally paid to it. The action of medicines is most rapidly produced when the drugs are injected under the skin, or into the muscles, blood, or spinal fluid, and next when they are taken upon an empty stomach. The rapid action of a purgative taken before breakfast, after the night's fast, is well known. On the other hand, medicines which have an irritating effect on the stomach, as arsenic or iodine, for instance, should be given soon after a meal. Iron appears to act best when dissolved by the acid of the gastric juice, and is therefore best given during active digestion. Cod-liver oil also agrees best if taken with, or soon after, a meal. The action of acids in checking the secretion of the gastric juice is taken advantage of by administering them a short time before meals, in those forms of dyspepsia accompanied with excessive secretion of acid gastric juice; whereas, if the contrary effects are desired, alkalies which excite the secretion of the gastric juice should be given

before food. The effects of valuable medicines are thus clearly influenced by their time of administration. In order that remedies may have the desired action, it is necessary that they be in a state of solution, or at least in a condition to render them easy of assimilation. Medicines which are insoluble in water, may be readily dissolved in the gastric juice; others again in the alkaline secretions of the pancreas and intestinal fluids. Minute division or the presence of fatty bodies also assists the rapid absorption and action of drugs. Care must therefore be taken in using solutions of corrosive sublimate to the head or other parts of the body, as it has been found that poisonous effects have been rapidly produced when a solution has been allowed to evaporate spontaneously on the skin.

The question of incompatibility is dealt with in the appendix, in which also tables of weights and measures in the Imperial and Metric Standards, and a table of approximate conversion from one to the other, will be found. A brief vocabulary of Latin words and phrases in common use in pharmacy, as well as tables of numerals and of the weights and measures translated into that language, are also appended.

ALIMENTARY DISEASES.

Acidity—

1

R Acidi Hydrochlorici Diluti	℥iss	5.3 cc.
Tincturæ Nucis Vomice	℥i	3.5 cc.
Infusi Gentianæ Compositi ad	℥vi	170 cc.
Fiat mistura. Misc.		
Sig. Capiat unciam ter in die ante cibos.		

2

R Sodii Bicarbonatis	gr. x	.65 gm.
Tincturæ Rhei Compositæ	℥xx	1.2 gm.
Tincturæ Camphoræ Compositæ	℥i	3.5 cc.
Aquæ Anethi ad	℥iss	45 cc.
Fiat haustus. Mitte vi tales. Solve et misce.		
Sig.—Capiendus si opus sit.		

3

R Ammonii Carbonatis	gr. v	.32 gm.
Potassii Chloratis	gr. x	.65 gm.
Tincturæ Opii	℥x.	.6 cc.
Infusi Lupuli ad	℥i	30 cc.
Fiat haustus. Mitte viii tales. Solve et misce.		
Sig.—Sumat sextis horis post cibum. (If painful.)		

4

R Bismuthi Carbonatis	℥i	4 gm.
Pulveris Acaciæ	℥iss	6 gm.
Creosoti	℥vi	.35 cc.
Spiritus Ammoniac Aromatici	℥ii	7 cc.
Syrupi Zingiberis	℥i	30 cc.
Aquæ ad	℥viii	230 cc.
Fiat mistura. Misc.		

Sig. Habeat cochlearia magna duo bis terve in die. P.P.A.

Aphthæ or Stomatitis -

5

R Boracis	℥i	4 gm.
Glycerini	℥ii	60 cc.
Aquæ ad	℥vi	170 cc.

Fiat collutorium. Solve et misce.

Sig.—Utendum sæpe.

6

R Sodii Sulphitis	gr. xl	2.6 gm.
Aquæ ad	℥i	30 cc.

Solve ut fiat applicatio.

Sig.—Apply with a small sponge or brush to the spots

7

R Potassii Chloratis	℥iii	12 gm.
Glycerini	℥i	30 cc.
Aquæ ad	℥xii	340 cc.

Fiat gargarisma. Solve et misce.

Sig.—Utendum sæpe.

Colic (intestinal)—

8

R Olei Ricini	℥iv.	15 cc.
Extracti Glycyrrhizæ Liquid	℥i	3.5 cc.
Tincturæ Opii	℥xx	1.2 cc.
Mucilaginis Acaciæ	℥iv	15 cc.
Aquæ Menthæ Piperitæ ad	℥iss	45 cc.

Fiat haustus. Misce.

Sig.—Statim sumendus. P. P. A.

9

R Tinct. Cardamoni Com-		
positæ	℥jv	15 cc.
Acidi Hydrocyanici Diluti	℥xl	2.4 cc.
Spiritus Ammoniz Aromatici	℥ij	7 cc.
Tinct. Zingiberis	℥iij	10.5 cc.
Aquæ Chloroformi ad	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Cochleare magnum pro re nata sumendum.

Constipation (chronic)—

10

R Hydrargyri Subchloridi - gr. ii .13 gm.
Pulveris Rhei Radicis - gr. v .32 gm.

Fiat pulvis. Mitte duodecim tales. Misce.

Sig.—Capiat unum horā somni, omni secundā nocte.

11

R Aloini - - - - -
Podophylli Resinæ - āā gr. ss .032 gm.

Extracti Nucis Vomicae - gr. 4 .016 gm.

Extracti Gentianæ - - q.s.

Fiat pilula. Mitte viginti tales. Misce.

Sig.—Capiat unam ter in die post cibos.

12

R Extracti Cascarae Sagradae

Liquidi - - - - - 5ii 60 cc.

Tincturae Nucis Vomicae - 5iii 10.5 cc.

Glycerini.

Aquae partes aequales ad 5vi 170 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare minimum omni mane nocteque.

13

R Extracti Aloes Barbadensis gr. i .065 gm.

Extracti Nucis Vomicae.

Extracti Belladonnae Viridis āā gr. ss .032 gm.

Fiat pilula. Mitte xxiv. tales. Misce.

Sig.—Capiat unam omni die ante prandium.

14

R Magnesii Sulphatis - - gr. cxx 8 gm

Mannae - - - - - gr. clx 10 gm.

Tincturae Sennae Co. - - 5ij 7 cc.

Infusi Sennae ad - - 5jss 45 cc.

Fiat haustus. Solve et misce.

Sig.—Sumat mane primo si opus sit.

(Similar to the common black draught.)

15

R Sodæ Tartaratæ	-	5jv	15 gm.
Tinct. Sennæ Co.	-	5ss	15 cc.
Syrupi Zingiberis	-	5ij	7 cc.
Aquæ ad	-	5jss	15 cc.

Fiat haustus. Solve et Misce.

Sig.—Si opus sit sum.

(More pleasant than the common black draught.)

16

R. Pulveris Rhei	-	-	-
Extracti Aloes Barbadosensis	āā	5ss	.2 gm.
Extracti Belladonnæ	-	gr. ij	.2 gm.
Olei Caryophylli	-	℥ij	.18 cc.

Divide into twenty pills. Misce.

Sig.—One night and morning.

17

R Extracti Aloes Barbadosensis	gr. ij	.2 gm.
Extracti Nucis Vomicae	-	-
Mastiche	-	āā gr. ss .032 gm.

Fiat pilula. Mitte xij tales. Misce.

Sig.—One pill night and morning.

18

R Podophylli Resinæ	-	gr. j	.065 gm.
Pilulae Rhei Compositæ	-	gr. x	.65 gm.
Extracti Hyoscyami	-	gr. jv	.26 gm.

Divide into four pills. Misce.

Sig.—One every third night.

(Sir G. Burrows).

19

R Podophylli Resinæ	-	gr. jv	.26 gm.
Pulveris Ipecacuanhæ	-	gr. vj	.4 gm.
Extracti Gentianæ	-	q.s.	-

Divide in pilulas duodecem. Misce.

Sig.—Una omni nocte sumenda.

20

R Magnesii Sulphatis . . . ̄j-ij 30-60 cc.
 Acidi Sulphurici Aromatici ̄jss 5.3 cc.
 Syrupi Zingiberis . . . ̄j 30 cc.
 Aquæ Chloroformi ad . . . ̄vj 170 cc.
 Fiat mistura. Solve et misce.
 Sig. Take ̄j every two or three hours.

21

R Extracti Cascaræ Sagradæ gr. iii .2 gm.
 Extracti Belladonnæ
 Extracti Nucis Vomice ̄ā gr. ̄ ̄.016 gm.
 Fiat pilula. Mitte xii tales. Misce.
 Sig.—Capiat unam bis in die.

22

R Magnesii Sulphatis . . . ̄j-ij 30-60 gm.
 Magnesii Carbonatis . . . ̄ss 15 gm.
 Sodii Bicarbonatis . . . ̄j 4 gm.
 Aquæ Menthæ Piperitæ ad ̄vj 170 cc.
 Fiat mistura. Solve et misce.
 Sig.—Take ̄j every two or three hours. P.P.A.

Constipation (acute)—

23

R Pilulæ Cambogiæ Composi-
 tæ
 Pilulæ Asafetidæ . . . ̄ā gr. v. .32 gm.
 Divide in pilulas duas. Mitte xii tales. Misce.
 Sig.—1 or 2 to be taken every night if necessary.
 (A good Purgative, acting chiefly on the Small
 Intestines.)

24

R Pilulæ Colocynthidis Com-
 positæ
 Pilulæ Hydrargyri . . . ̄ā gr. ijss .16 gm.
 Fiat pilula. Mitte xii tales. Misce.
 Sig.—Una nocte pro re nata sumenda.

25

R Olei Crotonis	-	-	-	℥½-1	.03-.06 cc.
Olei Caryophylli	-	-	-	℥j	.06 cc.
Micæ panis	-	-	-	q.s.	

Fiat pilula. Mitte duas tales. Misce.

Sig.—Statim una sumenda, et horis dualibus repetenda, si opus sit.

In Lead Colic.

26

Elaterii	-	-	-	gr. jss	.1 gm.
Pulveris Capsici	-	-	-	gr. jx	.6 gm.
Hydrargyri Subchloridi	-	-	-	gr. xij	.8 gm.
Extracti Hyoscyami	-	-	-	gr. xvij	1.2 gm.

Divide in pilulas viginti. Misce.

Sig. Sumat j vel ij si opus sit.

(An active purgative in lead colic. The Capsicum prevents the nausea produced by Elaterium).

Diarrhœa—

27

R Tinct. Catechu	-	-	-	ʒiij	10.5 cc.
Pulveris Cretæ Aromatici	-	-	-	ʒjss	6 gm.
Tinct. Opii	-	-	-	ʒss-j	1.8-3.5 cc.
Misturæ Cretæ ad	-	-	-	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ʒj post singulas liquidas sedes. P.P.A.

In mild cases.

28

R Extracti Bebe Liquidi	-	-	-	ʒij	7 cc.
Decocti Hordei	-	-	-	ʒjss	45 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Sumat post singulas sedes liquidas.

In simple cases.

29

R Tinct. Opii	-	-	3j	3.5 cc.
Olei Cajuputi	-	-	℥xij	.7 cc.
Tinct. Rhei	-	-	℥ss	15 cc.
Misturæ Cretæ ad	-	-	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Cochlearia magna duo post sedes liquidas sumenda.

In painful cases.

30

R Olei Ricini	-	℥vi	21 cc.
Spiritus Menthæ Piperitæ	-	℥x	.6 cc.
Spiritus Frumenti ad	-	℥iss	45 cc.

Fiat haustus. Misce.

Sig.—Statim sumendus.

In irritative cases.

31

R Bismuthi Carbonatis	-	-	-
Magnesii Carb. Levis	-	āā gr. xv	1 gm.
Mucilaginis Acaciæ	-	3i	3.5 cc.
Aquæ Anethi ad	-	℥j	30 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Sumendus ter in die si opus sit. P. P. A.

In dyspeptic cases.

32

R Cupri Sulphatis	-	-	-
Pulveris Opii	-	āā gr. ss	.032 gm.
Confectionis Rosæ	-	q.s.	-

Misce ut fiat pilula. Mitte xii tales.

Sig.—Una ter in die sumenda.

In chronic cases.

33

R Argenti Nitratis	-	-	gr. $\frac{1}{2}$.016-.032 gm.
Aque Destillatae	-	-	℥ij	60 cc.
Pulveris Acaciae	-	-	gr. xl	2.6 gm.
Sacchari Albi	-	-	℥ij	8 gm.

Fiat mistura. Misce.

Sig.—Sumat 3j secunda vel tertiâ quâque
horâ. P.P.A.

In chronic cases.

Dyspepsia—

34

R Pepsini	-	-	gr. lx	4 gm.
Creosoti	-	-	℥x	.6 cc.
Bismuthi Carbonatis	-	-	gr. lx	4 gm.

Fiat pulvis. Divide into thirty equal parts. Misce.
Sig.—Give one in a gelatine capsule thrice a day.
With flatulence.

35

R Pepsini Porci	-	-	gr. xxxij	2.1 gm.
Extracti Aloes Barbadosensis	-	-	gr. viij	.52 gm.
Glycerini	-	-	q.s.	

Fiat massa et divide in pilulas octo. Misce.
Sig.—Capiat unam nocte maneque.
With constipation.

36

R Pil. Colocynthis et Hyoscyami	-	-		
Fellis Bovini Purificati	-	-		
Extracti Lupuli	-	-	ââ gr xx	1.3 gm.

Misce et divide in pilulas viginti.
Sig.—One three hours after dinner.
In sluggish liver

37

R Acidi Nitro-Hydrochlorici

Diluti	-	-	-	℥iiss	5.3 cc.
Tincturæ Nucis Vomicae	-	-	-	℥i	3.5 cc.
Tincturæ Aurantii	-	-	-	℥iv	15 cc.
Infusi Calumbæ ad	-	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ℥iv ter in die ante cibos ex aquâ.

As a tonic.

38

R Acidi Nitro-hydrochlorici

Diluti	-	-	-	℥iii	10.5 cc.
Tincturæ Nucis Vomicae	-	-	-	℥iiss	5.3 cc.
Tincturæ Calumbæ	-	-	-	℥vi	21 cc.
Infusi Gentianæ Compositi	-	-	-	-	-
ad	-	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat coch. medium ex aquâ ante cibos.

As a tonic.

39

R Glycerini Pepsinæ	-	-	-	℥iiss	75 cc.
Tincturæ Chiratae	-	-	-	℥iv	15 cc.
Infusi Aurantii Compositi ad	-	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ℥ii ex aquâ post cibos.

In acidity.

40

R Resinæ Podophylli	-	-	-	gr. ss	.032 gm.
Extracti Euonymi Sicci	-	-	-	gr. iss	.1 gm.
Extracti Hyoscyami	-	-	-	q.s.	

Fiat pilula. Mitte xxiv. tales Misce.

Sig.—Capiat unam omni nocte.

In hepatic cases.

41

R Bismuthi Carbonatis	-	̄iv	15 gm.
Sodii Bicarbonatis.			
Magnesii Carb. levis	āā	̄iii	12 gm.
Pulveris Tragacanthæ	-	gr. xx	1.3 gm.
Liquoris Morphinae Hydro-			
chloridi	-	̄ii	7 cc.
Spiritus Chloroformi	-	̄iii	10.5 cc.
Aquæ ad	-	̄x	280 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ̄iv ter in die ante cibos. P.P.A.

In irritative cases.

42

R Potassii Bicarbonatis	-	̄i	4 gm.
Tinct. Nucis Vomicae	-	̄i	3.5 cc.
Infusi Gentianæ Co. ad	-	̄vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ̄j ter in die ante cibos.

In acidity.

43

R Ammonii Chloridi	-	̄ij	8 gm.
Tinct. Nucis Vomicae	-		
Tinct. Zingiberis	-	āā ̄j	3.5 cc.
Tinct. Gentianæ Compositæ	̄ss		15 cc.
Aquæ Chloroformi ad	-	̄vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ̄ss ter in die ante cibos.

In acidity.

44

R Sodii Bicarbonatis	-	̄jss	6 gm.
Spiritus Ammonia Aromatici	̄ij		7 cc.
Tinct. Gentianæ Compositæ.			
Syrupi Aurantii	-	āā ̄ss	15 cc.
Infusi Gentianæ Co. ad	-	̄vii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Habeat ̄j ter in die ante cibos.

In acidity.

Flatulence—

45

R Acidi Carbolici (cryst) - gr. i .065 gm.
 Pilulæ Aloes et Asafetidæ - gr. iii .2 gm.
 Fiat pilula. Mitte xxiv tales. Misce.
 Sig.—Capiat unam mane nocteque.

46

R Spiritūs Ammoniaë Aro-
 matici ʒiii 10.5 cc.
 Spiritūs Cajuputi ʒiiss 5.3 cc.
 Spiritūs Chloroformi ʒii 7 cc.
 Infusi Aurantii Compositi
 ad ʒvi 170 cc.
 Fiat mistura. Misce.
 Sig.—Capiat ʒiv si opus sit.

47

R Magnesii Carbonatis Levis - ʒii 8 gm.
 Spiritūs Menthaë Piperitæ - ʒi 3.5 cc.
 Spiritūs Chloroformi - ʒii 7 cc.
 Syrupi Zingiberis - ʒi 30 cc.
 Aquæ Anethi ad - ʒiii 90 cc.
 Fiat mistura. Misce.
 Sig.—Capiat ʒii dolore urgenti ex aquā. P.P.A.

48

R Salolis
 Bismuthi Salicylatis.
 Sodii Bicarbonatis - - - āā gr. v .32 gm.
 Fiat pulvis. Mitte xii tales. Misce.
 Sig.—Habeat unum bis terve in die ex lacte.

49

R Spiritūs Aetheris - - - ʒiij 10.5 cc.
 Spiritūs Chloroformi - - - ʒiij 10.5 cc.
 Tinct. Cardamomi Compositæ ʒvj 21 cc.
 Tinct. Rhei - - - ʒiij 10.5 cc.
 Aquæ Menthaë Piperitæ ad ʒvj 170 cc.
 Fiat mistura. Misce.
 Sig.—Sumat ʒss pro re nata.

Gall Stones—

50

R Tincture Belladonnæ	-	ʒi	3.5 cc.
Syrupi Chloralis	-	ʒvi	21 cc.
Aquæ ad.	-	ʒvi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒiv. tertiis horis.

Gastralgia

51

R Liq. Bismuthi et Ammonii			
Citratæ	-	ʒvi	21 cc.
Acidi Hydrocyanici Diluti	-	℥xv	.9 cc.
Tincturæ Nucis Vomice	-	ʒi	3.5 cc.
Tincturæ Aurantii	-	ʒiss	5.3 cc.
Syrupi Simplicis	-	ʒi	30 cc.
Aquæ Chloroformi ad	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat cochleare magnum ex aquâ quartis horis post cibum. P.P.A.

52

R. Liq. Bismuthi et Ammonii			
Citratæ	-	ʒi	3.5 cc.
Syrupi Aurantii	-	ʒiij	10.5 cc.
Infusi Chiratae ad	-	ʒjss	45 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Sumendus ter in die post cibum.

Gastric Ulcer—

53

R Bismuthi Subnitratæ	-	ʒiv	15 gm.
Acidi Hydrocyanici Diluti	ʒi		3.5 cc.
Liquoris Morphinae Hydroch.	ʒiii		10.5 cc.
Spiritus Chloroformi	-	ʒiii	10.5 cc.
Aquæ ad	-	ʒx	280 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒiv ter in die post cibos. P.P.A.

54

R Liquoris Arsenicalis - ℥j-vel ij .06-.12 cc.
 Aquæ ad - - - - - ̄ss 15 cc.
 Fiat haustus. Mitte xii tales. Misce.
 Sig. — Capiendus statim post cibum bis in die.

Hæmatemesis —

55

R Creosoti - - - - - ℥vj .35 cc.
 Olei Terebinthinæ - - - ̄ij 7 cc.
 Syrupi Aurantii - - - ̄vj 21 cc.
 Mucilaginis Acaciæ ad - ̄vj 170 cc.
 Fiat mistura. Misce.
 Sig. Sumat ̄j quartâquâque horâ. P.P.A.

56

R Tinct. Ferri Perchloridi - ℥xv .9 cc.
 Acidi Hydrochlorici Diluti - ℥x .6 cc.
 Olei Menthæ Piperitæ - ℥j .06 cc.
 Infusi Quassiae ad - - - ̄jss 45 cc.
 Fiat haustus. Mitte vi tales. Misce.
 Sig. — Habeat ter in die vel si opus sit.

57

R Extracti Ergotæ Liquidî - ℥xl 2.4 cc.
 Acidi Gallici - - - - - gr. x .65 gm.
 Aquæ Carui ad - - - - - ̄j 30 cc.
 Fiat haustus. Mitte viii tales. Misce.
 Sig. — Sumendus ter in die si opus sit.

58

R Pilulæ Plumbi cum Opio - gr. iv .26 gm.
 Mitte xii tales.
 Sig. Capiat unam quartis horis.

59

R Calcii Chloridi	-	-	-	℥ii	8 gm.
Tincturæ Aconiti	-	-	-	℥xii	.7 cc.
Acidî Hydrocyanici Diluti	-	-	-	℥xii	.7 cc.
Aquæ Chloroformi ad	-	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ̄ss tertiis horis si opus sit. P.P.A.

60

R Terebeni	-	-	-	℥x	.6 cc.
Mucilaginis Tragacanthæ.					
Glycerini	-	-	-	āā ̄i	3.5 cc.
Aquæ Cinnamomi ad	-	-	-	̄i	30 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Repetatur sextis horis si opus sit. P.P.A.

61

R Aluminis	-	-	-	̄j	4 gm.
Syrupi Rhæados	-	-	-	̄vj	21 cc.
Infusi Rosæ Acidî ad	-	-	-	̄viij	230 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ̄j ter in die.

Hæmorrhoids—

62

R Extracti Hæmamelidis Liq.	̄ss	15 cc.
Lanolini.		
Paraffini Mollis	-	āā ̄ij 8 gm.

Fiat unguentum. Misce.

Sig.—Applicandum ad anum post singulas sedes.

63

R Sulphuris Sublimati.		
Acidî Potassii Tartratis	āā ̄i	30 gm.

Fiat pulvis. Misce.

Sig.—Sum. ̄i ex lacte omni mane ante jentaculum.

64

R Confectionis Sennæ	-	-	
Sulphuris Sublimati	-	āā 5j	30 gm.
Pulveris Jalapæ	-	- 5j	4 gm.
Copaibæ	-	- 5jv	15 cc.
Pulveris Zingiberis	-	- 3ss	2 gm.
Potassii Tartratis Acidi	-	- 5jv	15 gm.
Syrupi Zingiberis	-	- q.s.	
Fiat electuarium. Misce.			
Sig.—Capiat 5i nocte maneque.			

65

R Confectionis Sulphuris	-	5ij	60 gm.
Extracti Taraxaci	-	- 5j	30 gm.
Misce.			
Sig.—A tea-spoonful in the morning.			

66

R Cocainæ Hydrochloridi	-	gr. xx	1.3 gm.
Morphinæ Hydrochloridi	-	gr. v	.32 gm.
Atropinæ Sulphatis	-	gr. jv	.26 gm.
Pulveris Acidi Tannici	-	gr. xx	1.3 gm.
Vaselini	-	- 3ss	15 gm.
Olei Rosæ	-	- q.s.	
Fiat ung. Misce.			

Sig.—Apply after each evacuation of the bowels.

67

R Chrysarobini	-	- gr. xv	1 gm.
Iodoformi	-	- gr. vj	.4 gm.
Extracti Belladonnæ	-	- gr. xij	.8 gm.
Vaselini	-	- 3vj	22 gm.

Fiat ung. Misce.

Sig.—Apply several times daily to the swellings, previously washed with a solution of carbolic acid.

68

R Unguenti Resinae	-	-	-
Unguenti Gallæ c Opio	aa	ʒvi	22 gm.
Vasellini ad	-	ʒij	60 gm.

Fiat ung. Misce.

Sig. Applicandum nocte maneque.

Sickness

69

R Potassii Bicarbonatis	-	ʒij	8 gm.
Spiritus Ammonia Aromatici	ʒij		7 cc.
Syrupi Aurantii	-	ʒj	30 cc.
Aquæ al	-	ʒvj	170 cc.

Solve et misce.

R Acidi Citrici	-	ʒiss	6 gm.
Aquæ ad	-	ʒvj	170 cc.

Fiat mistura. Solve.

Sig. Two table-spoonfuls of each to be mixed together and drank during effervescence every four hours.

70

R Extracti Opii	-	gr. j	.065 gm.
Pulveris Capsici	-	gr. ij	.13 gm.
Extracti Hyoscyami	-	gr. jv.	.26 gm.

Fiat massa. Divide in pilulas duas. Misce.

Sig.—Take one at bed-time.

Sea Sickness—

71

R Antipyriini	-	gr. lxxv	5 gm.
Cocainæ Hydrochloridi	-	gr. jss	.1 gm.
Caffeinæ	-	gr. jv	.26 gm.
Strychnina Sulphatis	-	gr. ʒss	.0022 gm.
Spiritus Vini Gallici	-	ʒiiss	9 cc.
Aquæ Destillatæ ad	-	ʒiij	90 cc.

Fiat mistura. Solve et misce.

Sig.—A table-spoonful before embarking, and others during the day, or three times during twenty-four hours.

Tape Worms—

72

R Extracti Filicis Liquidi	-	-	-
Syrupi Zingiberis	-	āā ʒi	3.5 cc.
Mucilaginis Acaciae Recentis	ʒss	-	5.3 cc.
Aquæ Cinnamomi ad	-	ʒi	30 cc.
Fiat haustus. Misce.			

Sig.—Sumendus mane alvo jejuna. P.P.A.

Note.—Male-Fern should always be followed in ten hours or so after administration by a dose of saline purgative, and not castor oil, as the latter renders felicie acid soluble, and causes it to be absorbed.

73

R Kamala	-	-	ʒi	4 gm.
Syrupi	-	-	ʒss	15 cc.
Pulveris Tragacanthæ	-	gr. vi	-	.4 gm.
Aquæ ad	-	-	ʒjss	45 cc.
Fiat haustus. Misce.				

Sig.—To be taken fasting and followed by a purge.
P.P.A.

74

R Cusso	-	-	ʒss	15 gm.
Mellis Depurati	-	-	ʒj	30 cc.
Fiat haustus. Misce.				

Sig.—Take half at once and the rest in six hours.

Thread Worms—(See p. 126-7.)

75

R Tinct. Ferri Perchloridi	-	ʒss	15 cc.
Aquæ ad	-	ʒviij	230 cc.
Misce.			

Sig.—Give as an enema.

Tympanitis—

76

R Olei Terebinthinæ	.	.	3j	3.5 cc.
Olei Amygdalæ Dulcis	.	.	℥ss	15 cc.
Tinct. Opii	.	.	℥ij	7 cc.
Mucilaginis Acaciæ	.	.	℥v	18 cc.
Aquæ Laurocerasi	.	.	℥ss	15 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare minimum tertiis vel sextis horis. P.P.A.

Vomiting (*to produce*)—

77

R Vini Antimonialis	.	.	℥ii	7 cc.
Vini Ipecacuanhæ	.	.	℥iii	10.5 cc.
Syrupi Scillæ ad	.	.	℥i	30 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare parvum quaque horæ parte quarta donec vomitus supervenerit.

78

R Cupri Sulphatis	.	.	gr. viii	.52 gm.
Aquæ	.	.	℥iij	90 cc.

Fiat haustus. Mitte duo tales. Solve.

Sig.—To be given at once, and repeated, if necessary, in half-an-hour.

79

R Zinci Sulphatis	.	.	gr. xx	1.3 gm.
Aquæ ad	.	.	℥iij	90 cc.

Fiat haustus. Mitte duo tales. Solve.

Sig.—Statim sumendus. Repetendus post horam dimidiam si opus sit.

BLOOD DISEASES.

Anæmia—

80

R Ferri et Ammonii Citratis	-	̄iii	12 gm.
Liq. Arsenici Hydrochlorici		̄i	3.5 cc.
Tincturæ Nucis Vomicae	-	̄ii	7 cc.
Spiritus Chloroformi	-	̄ii	7 cc.
Aquæ ad	- - -	̄vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat cochleare medium ex aquâ ter in die post cibos.

81

R Ferri Sulphatis	- - -	gr. ijss	.16 gm.
Potassii Carbonatis	- - -	gr. jss	.1 gm.
Sacchari	- - -	gr. j	.065 gm.
Tragacanthæ	- - -	gr. ʒ	.01 gm.

Fiat pilula. Mitte quinquaginta tales. Misce.

Sig.—Sumendæ duæ ter in die post cibos

(Similar to Bland's Pill).

82

R Tinct. Ferri Perchloridi	-	̄ij	7 cc.
Liq. Strychninae Hydroch-			
loridi	- - -	̄j	3.5 cc.
Glycerini	- - -	̄ss	15 cc.
Infusi Quassiaæ ad	- - -	̄vj	170 cc.

Fiat mistura. Misce.

Sig.—Habeat ̄ss ter in die post cibum ex aquâ.

Anæmia (Pernicious).—

83

R	Liquoris Arsenicalis	-	-	℥xxxvj	2.12 cc.
	Tinct. Lavandulæ Co.	-	-	ʒj	3.5 cc.
	Aquæ ad	-	-	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ʒss ter in die post cibos ex aquâ.

In early stage.

84

R	Liquoris Arsenicalis	-	-	ʒiss	5.3 cc.
	Tincturæ Cardamomi Com-				
	positæ	-	-	ʒvi	22 cc.
	Aquæ Menthæ Piperitæ ad	-	-	ʒviii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat coch. amplum ter in die post cibos ex aquâ.

*In later stage.***Chlorosis—**

85

R	Ferri et Ammonii Citratis	-	-	ʒii	8 gm.
	Liquoris Arsenici Hydroch-				
	lorici	-	-	ʒi	3.5 cc.
	Magnesi Sulphatis	-	-	ʒi	30 gm.
	Glycerini	-	-	ʒvi	21 cc.
	Aquæ ad	-	-	ʒviii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. magnum ter in die post cibos ex aquâ.

Hæmophilia—

86

R	Potassii Chloratis	-	-	ʒii	8 gm.
	Tincturæ Ferri Perchloridi	-	-	ʒiii	10.5 cc.
	Syrupi Limonis	-	-	ʒi	30 cc.
	Aquæ Chloroformi ad	-	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. magnum ter in die post cibos.

Purpura —

S7

R Tincturæ Ferri Perchloridi	̄iii	10.5 cc.
Liquoris Arsenici Hydroch-		
lorici	̄i	3.5 cc.
Glycerini	̄vi	21 cc.
Aquæ ad	̄vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat coch. magnum ter in die post cibos ex aquâ.

CIRCULATORY DISEASES.

Aneurysm—

88

R Potassii Iodidi	̄v	18 gm.
Calcii Chloridi	̄iv	15 gm.
Extracti Glycyrrhizæ Liquidi	̄i	30 cc.
Aquæ ad	̄x	280 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ̄iv ter in die post cibos ex aquâ.

S9

R Potassii Iodidi	gr. xv	1.3 gm.
Aquæ Chloroformi ad	̄i	30 cc.

Fiat haustus. Mitte viii tales. Solve.

Sig.—A draught to be taken in water every four hours until four have been taken, and afterwards twice a day after food.

Angina Pectoris—

90

R Liquoris Trinitrini	-	-	℥ii	.12 cc.
Tincturæ Cardamomi Com-				
positæ	-	-	℥ss	1.8 cc.
Aquæ ad	-	-	℥i	30 cc.

Fiat haustus. Mitte sex tales. Misce.

Sig.—Statim sumendus et quartis horis repetendus
si opus sit.*During the spasm.*

91

R Amylis Nitritis	-	-	℥iii	.18 cc.
Pone in capsulam vitream.				Mitte vi tales.

Sig.—To be broken in a handkerchief and the
vapour inhaled. Repeat if necessary in an hour.*During the spasm.*

92

R Amylis Nitritis	-	-	℥xii	.7 cc.
Ethylic Alcohol (Sp. gr '834)			℥v	18 cc.
Glycerini ad	-	-	℥jss	45 cc.

Fiat mistura. Misce.

Sig.—Habeat cochleare parvum ex fervente aquâ
tertiis horis si opus sit.*Between spasms.*

93

R Potassii Iodidi	-	-	℥ii	8 gm.
Tincturæ Strophanthi	-	-	℥iii	10.5 cc.
Liquoris Arsenicalis	-	-	℥iss	5.3 cc.
Aquæ Chloroformi ad	-	-	℥vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. medium ex aquâ ter in die
post cibos.*In chronic cases.*

94

R Spiritus Ætheris Co. -	-	5j	3.5 cc.
Liquoris Opii Sedativi -	-	℥x	.6 cc.
Aquæ Camphoræ ad -	-	℥ss	15 cc.

Fiat haustus. Mitte xii tales. Misce.

Sig. Capiat pro re nata.

In chronic cases.

Dropsy (Cardiac)—

95

R Tinct. Scillæ -	-	-	5ij	7 cc.
Tinct. Camphoræ Com-	-	-	-	-
posite -	-	-	5vj	21 cc.
Liquoris Ammonii Citratis -	-	-	℥ij	60 cc.
Infusi Scoparii ad -	-	-	℥viij	230 cc.

Fiat mistura. Misce.

Sig.—Sumat coch. magna duo ter in die.

96

R Potassii Acetatis -	-	-	℥ss	2 gm.
Aceti Scillæ -	-	-	℥ss	1.8 cc.
Spiritus Ætheris Nitrosi -	-	-	℥xx	1.2 cc.
Tinct. Digitalis -	-	-	℥v	.3 cc.
Infusi Scoparii ad -	-	-	℥jss	45 cc.

Fiat haustus. Mitte vi tales. Solve et misce.

Sig.—Ter in die sumendus.

97

R Potassii Acetatis -	-	-	℥jss	6 gm.
Spiritus Ætheris Nitrosi -	-	-	5ij	7 cc
Tinct. Digitalis -	-	-	-	-
Tinct. Scillæ -	-	-	ââ 5j	3.5 cc.
Infusi Digitalis Recentis ad	-	-	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat coch. largum ter in die. (Where there is diminished arterial pressure.)

98

R Pulveris Digitalis . . .
 Pulveris Scillæ āā gr. xij .8 gm.
 Extracti Taraxaci q.s.

Fiat massa. Divide into twelve pills. Misce.

Sig.—One to be taken twice daily.

As a Diuretic in Mitral, but dangerous in Aortic Disease (Dr Tanner).

Endocarditis—

99

R Sodii Salicylatis ʒiii. 12 gm.
 Sodii Bicarbonatis ʒv 18 gm.
 Tincturæ Nucis Vomicae ʒi 3.5 cc.
 Glycerini ʒvi 21 cc.
 Aquæ ad ʒvi 170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat cochleare amplum quartis horis ex aquâ.

In acute cases.

100

R Pulveris Digitalis
 Pulveris Scillæ
 Pilulæ Hydrargyri āā gr. i .065 gm.

Fiat pilula. Mitte xxiv tales. Misce.

Sig.—Capiat unam ter in die post cibos.

In chronic cases.

Heart Disease—

101

R Ferri et Quininae Citratis ʒjv 15 gm.
 Tinct. Nucis Vomicae ʒiij 10.5 cc.
 Syrupi Simplicis ad ʒvj 170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat coch. minima duo post cibos ex aquâ.

In irritable heart.

102

R Tinct. Digitalis	-	-	℥x	.6 cc.
Acidi Hydrocyanici Diluti	-	-	℥iij	.18 cc.
Tinct. Opii	-	-	℥v	.3 cc.
Aquæ Camphoræ ad	-	-	℥j	30 cc.

Fiat haustus. Mitte viii tales. Misce.

Sig.—Capiat ter in die. P. P. A.

In irritable heart.

103

R Tincturæ Digitalis	-	-	℥iss	5.3 cc.
Tincturæ Nucis Vomicae	-	-	℥i	3.5 cc.
Spiritus Ammoniaë Aromatici	-	-	℥iv	15 cc.
Aquæ Chloroformi ad	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat coch. magnum ex aquâ ter in die.

In weak heart.

104

R Tinct. Digitalis	-	-	℥x	.6 cc.
Ferri et Ammonii Citratis	-	-	-	-
Ammonii Carbonatis	-	-	āā gr. v	.32 gm.
Aquæ ad	-	-	℥j	30 cc.

Fiat haustus. Mitte viii tales. Solve et misce.

Sig.—Sumat ter in die ex aquâ post cibum.

In weak heart

105

R Pulveris Digitalis	-	-	gr. j	.065 gm.
Ferri Sulphatis Exsiccati	-	-	gr. ʒ	.016 gm.
Pulveris Capsici	-	-	gr. j	.065 gm.
Extracti Aloes	-	-	gr. ij	.12 gm.

Fiat pilula. Mitte xii tales. Misce.

Sig.—Habeat unam bis in die.

In weak heart.

Pericarditis—

106

R Sodii Salicylatis . . .	5v	18 gm.
Ammonii Carbonatis . . .	5ss	2 gm.
Tincturæ Nucis Vomicae . .	5iiss	5.3 cc.
Aquæ ad . . .	5viii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. plenum sextis horis.

CONSTITUTIONAL DISEASES.

Gout (acute)—

107

R Magnesii Sulphatis . . .	5jss	45 gm.
Magnesii Carbonatis Levis .	5ss	2 gm.
Vini Colchici . . .	5j	3.5 cc.
Aquæ Chloroformi ad . . .	5vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat coch. magna duo ter in die. P.P.A.

In acute cases.

108

R Potassii Bicarbonatis . . .	5vi	22 gm.
Potassii Citratis . . .	5iv	15 gm.
Vini Colchici . . .	5vi	21 cc.
Spiritus Chloroformi . . .	5iii	10.5 cc.
Aquæ ad . . .	5viii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. plenum quartis horis.

In acute cases.

109

R Potassii Citratis . . .	5ii	8 gm.
Tinct. Stramonii . . .	5j	3.5 cc.
Tinct. Colchici . . .	5ij	7 cc.
Tinct. Digitalis . . .	5i	3.5 cc.
Aquæ Menthæ Piperitæ ad .	5vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Coch. larga duo ter in die sumenda.

In asthmatic cases.

110

R Vini Colchici	-	-	-	
Tinct. Opii	-	-	āā ʒss	15 cc.
Sodii Carbonatis	-	-	ʒiv	15 gm.
Aquæ ad	-	-	ʒiij	90 cc.

Fiat lotio. Solve et misce.

Sig.—To be sprinkled on cloths wrung out of hot water, and applied to the joints.

When localised.

111

R Collodii Flexilis	-	-	-	
Ætheris	-	-	āā ʒv	18 cc.
Acidi Salicylici	-	-	ʒj	4 gm.
Morphinæ Hydrochlor.	-	-	ʒss	2 gm.

Fiat pig. Solve et misce.

Sig.—Pinge super partem dolentem.

When localised.

Gout (chronic)—

112

R Extracti Colchici	-	-	gr. iss	.1 gm.
Extracti Rhei	-	-	-	
Extracti Aloes Barbadosensis	āā	gr. vj		.4 gm.
Extracti Belladonnæ Alcoholici	gr. j			.065 gm.
Fiat massa.	Divide in pilulas sex. Misce.			

Sig.—One at night twice a week.

113

R Sodii Bicarb.	-	-	ʒj	4 gm.
Sodii Chloridi	-	-	gr. iv	.26 gm.
Sodii Sulphatis	-	-	gr. x	.65 gm.
Magnesii Sulphatis	-	-	gr. iij	.2 gm.
Ferri Sulphatis	-	-	gr. i	.065 gm.
Aquæ ad	-	-	ʒxx	568 cc.

Fiat mistura. Solve.

Sig.—Capiat cyathum amplum omni mane.
(An imitation of Vichy Water.)

114

R Potassii Iodidi	.	.	gr. xv	1 gm.
Vini Colchici	.	.	ʒjss	5.3 cc.
Tinct. Hyoscyami	.	.	ʒj	3.5 cc.
Infusi Calumbæ ad	.	.	ʒvj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat cochlearia magna duo ter in die ex aquâ post cibum.

115

R Lithii Carbonatis	.	.	gr. v	.32 gm.
---------------------	---	---	-------	---------

Fiat pulvis. Mitte xii tales.

Sig.—Take one in half a bottle of soda water thrice daily.

116

R Acidi Hydrochlorici Diluti	ʒjss	5.3 cc.
Acidi Hydrocyanici Diluti	℥xx	1.2 cc.
Infusi Chiratae ad	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Cochlearia magna duo ter in die sum.

In dyspeptic cases.

Rheumatism (acute)—

117

R Sodii Salicylatis	.	.	ʒiii	12 gm.
Potassii Bicarbonatis	.	.	ʒiv	15 gm.
Magnesii Sulphatis	.	.	ʒvi	22 gm.
Tincturæ Nucis Vomicae	.	.	ʒiss	5.3 cc.
Aquæ ad	.	.	ʒviii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. plenum tertiis horis ex aquâ.

118

R Sodii Salicylatis	.	.	ʒjss	6 gm.
Syrupi Simplicis	.	.	ʒj	30 cc.
Aquæ ad	.	.	ʒvj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Cochlearia magna duo duabus horis donec quartæ doses captæ sint ; postea quartis horis.

119

R Sodii Salicylatis	ʒij	4 gm.
Liquoris Ammonii Acetatis	ʒij	60 cc.
Aquæ Chloroformi ad	ʒvj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. magna duo quartâquâque horâ.

Rheumatism (chronic)—

120

R Potassii Iodidi	ʒij	8 gm.
Infusi Gent. Co.	ʒviij	230 cc.

Fiat mistura. Solve.

Sig.—Capiat coch. mag. duo ter in die ex aquâ
post cibum.

121

R Quininæ Sulphatis	ʒj	4 gm.
Liquoris Sodii Arsenatis	ʒiij	10.5 cc.
Acidi Sulphurici Dil.	ʒj	3.5 cc.
Tinct. Cinchonæ Co.	ʒij	60 cc.
Syrupi Zingiberis ad	ʒiij	90 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat cochleare parvum bis terve die
post cibos.*In debility.*

122

R Potassii Iodidi		
Ammonii Chloridi	ʒjss	6 gm.
Infus Lupuli ad	ʒvj	170 cc.

Fiat mistura. Solve.

Sig.—Sumat coch. largum ter in die ex aquâ
post cibum.

123

R Tinct. Guaiaci Ammoniatæ	̄iij	10.5 cc.
Tinct. Aconiti	̄lxx	1.2 cc.
Misturæ Amygdalæ ad	̄vj	170 cc.
Fiat mistura. Misce.		

Sig.—Sumat coch. mag. duo ter in die.

124

R Linimenti Belladonnæ	-	
Linimenti Opii	̄ā ̄i	30 cc.
Linimenti Terebinthinæ	̄ij	60 cc.
Fiat linimentum. Misce.		

Sig.—Infricetur bene affectis partibus.

When local.

125

R Olei Gaultheriæ	-	
Olei Olivæ	̄ā ̄ss	15 cc.
Fiat pigmentum. Misce.		

Sig.—Applicandum affectis partibus mane nocteque.

When local.

126

R Tincturæ Cimicifugæ	̄i	30 cc.
Tincturæ Belladonnæ	̄ii	7 cc.
Aquæ Chloroformi ad	̄viii	230 cc.
Fiat mistura. Misce.		

Sig.—Uncia sumenda quartis horis donec dolor mitigatus sit.

In lumbago.

127

R Linimenti Camphoræ Ammoniatæ		
Linimenti Belladonnæ	̄ā ̄jss	45 cc.
Fiat linimentum. Misce.		

Sig.—Perfricandum supra dorsum nocte maneque fortiter.

In lumbago

EAR, EYE, NOSE, AND THROAT DISEASES.

Earache—

128

R Tinct. Opii - - - - -
 Olei Amygdalæ - - - - - āā ñiv 15 cc.
 Fiant guttæ. Misce.

Sig. Una ad tres guttæ instillandæ in aurem si
 opus sit.

129

R Resorcini - - - - - gr. jss .1 gm.
 Cocainæ Hydrochloridi - gr. v .32 gm.
 Morphinae Hydrochloridi - gr. ¼ .01 gm.
 Aquæ Destillatæ ad - - - - - ñi 30 cc.

Fiant guttæ. Solve et misce.

Sig.—For external use by instillation.

Eye (inflammation)—

130

R Acidi Borici - - - - - ñss 2 gm.
 Aquæ Destillatæ - - - - - ñii 90 cc.

Fiat collyrium. Solve.

Sig.—Utendum tertiis vel quartis horis.

131

R Aluminis - - - - - gr. jv .26 gm.
 Aquæ ad - - - - - ñj 30 cc.

Fiant guttæ. Solve.

Sig.—Use two or three drops twice a day,
 or oftener if necessary.

132

R Argenti Nitratis	-	-	gr. ij	.13 gm.
Aquæ Destillatæ	-	-	ʒj	30 cc.
Fiant guttæ. Solve.				
Sig.—Utendæ more dicto.				

133

R Aluminis	-	-	gr xxjv	1.5 gm.
Atropinæ Sulphatis	-	-	gr. j	.065 gm.
Aquæ Destillatæ	-	-	ʒviiij	230 cc.
Fiat lotio. Solve.				
Sig. Utenda ut dictum.				

134

R Atropinæ Sulphatis	-	-	gr. i	.065 gm.
Aquæ Destillatæ	-	-	ʒss	15 cc.
Fiant guttæ. Solve.				
Sig.—Una vel duæ instillandæ si opus sit.				

135

R Cocainæ Hydrochloridi	-	-	gr. vi	.4 gm.
Boracis	-	-	-	-
Acidi Borici	-	-	ʒss	2 gm.
Aquæ Destillatæ ad	-	-	ʒiii	90 cc.
Fiant guttæ. Solve.				
Sig.—Duæ vel tres in oculos instillantur m. et n.				

Nose and Throat Inflammations—

136

R Mentholis	-	-	-	-
Camphoris	-	-	ʒss	2 gm.
Olei Cinnamomi	-	-	ʒv	.3 cc.
Paraffini Liquidi ad	-	-	ʒi	30 cc.
Fiat nebula. Solve et misce.				
Sig.—To be used with an atomiser as directed.				

137

R Eucalyptolis
Olei Gaultheriæ	.	.	āā ℥v	.3 cc.
Mentholis	.	.	gr. x.	.65 cc.
Paraffini Liquidi ad	.	.	ʒi	30 cc.

Fiat nebula. Solve et misce.

Sig.—To be used with an atomiser as directed.

138

R Cocainæ Hydrochloridi	.	gr. ii	.13 gm.
Olei Cinnamomi	.	℥v	.3 cc.
Mentholis	.	gr. xv	1 gm.
Paraffini Liquidi ad	.	ʒi	30 cc.

Fiat nebula. Solve et misce.

Sig.—To be used with an atomiser as directed.

Throat (relaxed)—

139

R Aluminis	.	.	ʒjss	6 gm.
Tinct. Capsici	.	.	ʒij	7 cc.
Infusi Rosæ Acidi ad	.	.	ʒviiij	230 cc.

Fiat gargarisma. Solve et misce.

Sig. Utendum pro re nata.

140

R Zinci Sulphatis	.	.	gr. xv	1 gm.
Syrupi Mori	.	.	ʒjv	15 cc.
Glycerini	.	.	ʒj	30 cc.
Infusi Krameriæ ad	.	.	ʒviiij	230 cc.

Fiat gargarisma. Solve et misce.

Sig.—Utendum sæpe.

141

R Creosoti	.	.	℥xx	1.2 cc.
Tinct. Lavandulæ Compositæ
Tinct. Myrrhæ	.	.	āā ʒjv	15 cc.
Glycerini	.	.	ʒi	30 cc.
Aquæ ad	.	.	ʒviiij	230 cc.

Fiat gargarisma. Misce.

Sig.—Utatur mane nocteque. P.P.A.

142

R Acidi Tannici	-	-	-	℥ss	15 gm.
Glycerini	-	-	-	℥jv	115 cc.

Fiat pigmentum. Solve.

Sig.—Applicandum gutturi ope penicilli, vel dilutum cum aquā pro gargarismate utendum.

Tonsillitis—

143

R Iodi	-	-	-	gr. vj	.4 gm.
Potassii Iodidi	-	-	-	gr. xij	.8 gm.
Mentholis	-	-	-	gr. x	.65 gm.
Glycerini ad	-	-	-	℥j	30 cc.

Fiat pigmentum. Solve.

Sig.—Applicet ad tonsillas more dicto. P.P.A.

144

R Sodii Salicylatis	-	-	-	℥ii	8 gm.
Tinct. Aconiti	-	-	-	℥ss	1.8 cc.
Syrupi Aurantii	-	-	-	℥j	30 cc.
Aquæ ad	-	-	-	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat coch. magnum ter in die.

145

R Potassii Chloratis	-	-	-	℥iii	12 gm.
Tincturæ Ferri Perchloridi	-	-	-	℥iv	15 cc.
Glycerini	-	-	-	-	-
Syrupi Limonis	-	-	-	āā ℥i	30 cc.
Aquæ ad	-	-	-	℥xii	340 cc.

Fiat gargarisma. Solve et misce.

Sig.—Two tablespoonfuls in an equal quantity of water to be gargled every four hours and afterwards swallowed.

146

R Tinct. Guaiaci Ammoniatæ ʒiij 10.5 cc.
 Tinct. Aconiti - - - ℥xv .9 cc.
 Misturæ Amygdalæ ad - ʒvj 170 cc.
 Fiat mistura. Misce.
 Sig.—Sumat coch. magna duo ter in die.

147

R Potassii Chloratis - - - gr. v .32 gm.
 Tinct. Ferri Perchloridi - - -
 Spiritus Chloroformi - āā ℥x .6 cc.
 Aquæ ad - - - ʒj 30 cc.
 Fiat haustus. Mitte viii tales. Solve et misce.
 Sig.—Sumendus quartâquâque horâ.

INFECTIOUS DISEASES.

Cholera—

148

R Acidi Sulphurici Diluti - ʒjss 5.3 cc.
 Tinct. Opii - - - ʒj 3.5 cc.
 Tinct. Cinnamomi - - ʒiij 10.5 cc.
 Syrupi - - - ʒss 15 cc.
 Aquæ Camphoræ ad - ʒvj 170 cc.
 Fiat mistura. Misce.
 Sig.—One ounce every time the bowels are moved.
 (In early stages.)

149

R Salolis - - - - - gr. v .32 gm.
 Spiritus Chloroformi - - ℥xx 1.2 cc.
 Mucilaginis Acaciæ - - ʒj 3.5 cc.
 Aquæ ad - - - ʒj 30 cc.
 Fiat haustus. Mitte vi tales. Solve et misce.
 Sig.—To be given every three or four hours.
 P. P. A.

150

R Tinct. Opii	-	-	-	3j	3.5 cc.
Tinct. Capsici	-	-	-	.	.
Spiritus Camphoræ	-	-	-	āā 3j	30 cc.
Chloroformi	-	-	-	3iij	10.5 cc.
Alcoholis Ethylicis 60% ad	-	-	-	3iv	115 cc.

Fiat mistura. Misce.

Sig.—Take ℥xx-xl in water every two or three hours.

151

R Acidi Sulphurosi	-	-	-	3i	30 cc.
Aquæ Chloroformi ad	-	-	-	3viii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare plenum ex aquæ cyatho medio vel succi lemonis recentis quartis horis.

In cholera nostras.

Diphtheria—

152

R Tinct. Ferri Perchloridi	-	-	-	3ss	1.8 cc.
Potassii Chloratis	-	-	-	3j	4 gm.
Syrupi	-	-	-	3ss	15 cc.
Aquæ ad	-	-	-	3vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat 3j quatuorquæ horâ.

153

R Potassii Chloratis (pulv.)	-	gr. x	.65 gm.
Acidi Hydrochlorici	-	3ss	1.8 cc.

Mix in a pint bottle and let the gas generate; cork the bottle for two minutes, and then add gradually, shaking after each addition, a pint of distilled water.

Sig.—Use frequently as a gargle, swallowing a little occasionally.

(Euchlorine gargle.)

154

R Acidi Lactici	-	-	-	℥i	3.5 cc.
Aquæ ad	-	-	-	℥ii	60 cc.

Fiat nebula. Misce.

Sig.—Applicanda sæpe ad fauces.

Dysentery—

155

R Liquoris Ferri Pernitratis	-	℥jss	5.3 cc.
Syrupi Simplicis	-	℥ss	15 cc.
Aquæ Anethi ad	-	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat cochleare magna duo ter in die
cum pilula sequente

R Extracti Hæmatoxyli	-	gr. v	.32 gm.
Spiritus Rectificati	-	q.s.	

Fiat pilula. Mitte vi tales. Misce.

156

R Decocti Hæmatoxyli	-	℥j	30 cc.
Acidi Nitrici Diluti	-	℥x	.6 cc.
Tinct. Opii	-	℥v	.3 cc.

Fiat haustus. Mitte viii tales. Misce.

Sig.—Capiat t. i. d. si opus sit.

157

R Pulveris Kino Compositi	-	gr. x	.65 gm.
Infusi Cascarillæ	-	℥vj	21 cc.
Syrupi Papaveris	-	℥j	3.5 cc.
Aquæ Cinnamomi ad	-	℥iss	45 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Capiendus bis terve in die. P.P.A.

In chronic cases.

158

R Emetinæ Hydrochloridi	-	gr. i	.65 gm.
Syrupi Aurantii	-	ʒi	30 cc.
Aquæ Chlorformi ad	-	ʒviii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒss to ʒi ter in die et dosin
minorem si vomitus supervenerit.*In amœbic dysentry.*

Erysipelas—

159

R Potassii Nitratis	-	gr. x	.65 gm.
Sodii Bicarbonatis	-	ʒss	2 gm.
Vini Antimonialis	-	ʒij	7 cc.
Syrupi Aurantii	-	ʒi	30 cc.
Spiritus Ætheris Nitrosi	-	ʒj	3.5 cc.
Aquæ ad	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Habeat cochlearia magna duo ter in die.

160

R Tincturæ Ferri Perchloridi	ʒiv	15 cc.
Potassii Chloratis	ʒi	4 gm.
Liquoris Ammonii Citratis	ʒiij	90 cc.
Aquæ Chloroformi ad	ʒviii	230 cc.

Fiat mistura. Misce.

Sig.—Sumat ʒss ad ʒi ter in die.

161

R Acidi Sulphurosi	-	-
Glycerini	-	āā ʒi 30 cc.

Fiat applicatio. Misce.

Sig.—Applicandum ad eutem integram ut dictum.

162

R Unguenti Ichthyolis	-	-
Vasellini	-	āā ʒss 15 gm.

Fiat unguentum. Misce.

Sig.—Applicandum affectis partibus.

Gonorrhœa—

163

R Magnesii Sulphatis	-	-	3ii	60 gm.
Tincturæ Hyoscyami	-	-	3vi	21 cc.
Tincturæ Belladonnæ	-	-	3iss	5.3 cc.
Infusi Buchu ad	-	-	3viii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat 3iv quater in die.

164

R Balsami Copaibæ	-	-	3vi	21 cc.
Pulveris Cubebæ	-	-	3i	4 gm.
Pulveris Tragacanthæ	-	-	3iii	12 gm.
Spiritus Æthieris Nitrosi	-	-	3iv	15 cc.
Aquæ Cinnamomi ad	-	-	3x	280 cc.

Fiat mistura. Misce. P.P.A.

Sig.—Capiat 3iv quater in die.

165

R Olei Copaibæ	-	-	3ij	7 cc.
Liquoris Potassæ	-	-	3j	3.5 cc.
Tinct. Lavandulæ Compositæ	-	-	3iij	10.5 cc.
Aquæ Camphoræ ad	-	-	3viij	230 cc.

Fiat mistura. Misce.

Sig.—Sumat 3j ter in die.

166

R Ichthyolis	-	-	3j	4 gm.
Aquæ ad	-	-	3viij	230 cc.

Fiat injectio. Solve.

Sig.—Half an ounce to be used thrice daily.

Hay Fever—

167

R Adrenalini Chloridi Solutionis	(1:1,000)	-	-	
Normalis Salini Solutionis	aa	3i		3.5 cc.

Fiat nebula. Misce.

Sig.—To be injected into the nose with an atomiser cautiously.

Influenza—

168

R Quininæ Sulphatis	-	-	̄ss	2 gm.
Acidi Citrici	-	-	̄iv	15 gm.
Infusi Aurantii Compositi ad			̄vi	170 cc.
Fiat mistura. Solve.				
Sig.—No. 1.				

R Ammonii Carbonatis	-	-	̄i	4 gm.
Potassii Bicarbonatis	-	-	̄v	18 gm.
Aquæ ad	-	-	̄vi	170 cc.
Fiat mistura. Solve.				

Sig.—One tablespoonful to be mixed with an equal quantity of No. 1, and taken every three hours.

Intermittent Fever—

169

R Beberinæ Sulphatis	-	-	̄ss	2 gm.
Syrupi	-	-	̄j	30 cc.
Acidi Sulphurici Aromatici			̄i	3.5 cc.
Tinct. Aurantii	-	-	̄j	30 cc.
Aquæ ad	-	-	̄vi	170 cc.
Fiat mistura. Misce.				
Sig.—Capiat ̄ss t.i.d.				

Leprosy—

170

R Olei Gynocardie (l.C.)	-	̄i	30 cc.
--------------------------	---	----	--------

Sig.—10 to 60 drops in capsules, cod-liver oil, or milk, thrice a day after meals. Also to be rubbed into the affected parts where the skin is not broken.

Malaria—

171

R Quininae Hydrobromidi . gr. x .6 gm.
Fiat pulvis. Mitte duodecim tales.

Sig.—One to three powders to be taken daily in lemon juice, or one given dissolved in distilled water as a hypodermic injection.

172

R Liquoris Arsenici Hydrochlorici ʒss 1.8 cc.
Tincturae Ferri Perchloridi ʒij 7 cc.
Aqua Chlorformi ad ʒviii 230 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒss vel ʒi ter in die post cibos ex equā.

173

R Quininae Hydrobromidi
Acidi
Urethane ʒā gr. x .65 gm.
Aqua destillata ad ʒii 7 cc.

Fiat injectio (hyp.). Solve et misce.

Sig.—5-10 minims to be injected into the muscles in severe æstivo-autumnal ague.

Plague—

174

R Tincturae Cinchonae Compositae ʒiv 15 cc.
Animonii Carbonatis ʒi 4 gm.
Tincturae Nucis Vomicae ʒiiss 5.3 cc.
Tincturae Digitalis ʒi 3.5 cc.
Aqua ad ʒvi 170 cc.

Fiat mistura. Solve et misce.

Sig.—Cochleare magnum ex aquā t.d.s.

Pyrexia—

175

R Succi Limonis	.	.	.	℥jv	115 cc.
Corticis Limonis Recentis	.	.	.	℥ss	15 cc.
Potassii Nitratis	.	.	.	℥j	4 gm.
Aquæ Ferventis	.	.	.	℥iij	1.7 litre

Fiat mistura. Solve et misce.

Sig.—Pro potu communi in febris capienda.

176

R Potassii Bitartratis	.	.	.	℥j	30 gm.
Olei Limonis	.	.	.	℥xv	.9 cc.
Sacchari Albi	.	.	.	℥iij	60 gm.
Aquæ Bullientis ad	.	.	.	℥iij	1 litre

Fiat mistura. Solve et misce.

Sig.—To be used as a drink freely.

177

R Potassii Acetatis	.	.	.	℥iv	15 gm.
Spiritus Ætheris Nitrosi	.	.	.	℥vi	21 cc.
Spiritus Chloroformi	.	.	.	℥ii	7 cc.
Aquæ ad	.	.	.	℥vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. mag. ex ℥i aquæ tertiis horis.

178

R Liquoris Ammonii Acetatis	.	.	.	℥iss	45 cc.
Vini Ipecacuanhæ	.	.	.	℥ij	7 cc.
Syrupi Tolutani	.	.	.	℥i	30 cc.
Tincturæ Cocci	.	.	.	℥i	3.5 cc.
Aquæ Chloroformi ad	.	.	.	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Cap. coch. magnum quartis horis ex aquâ.

179

R Potassii Nitratis	.	.	gr. xx	1.3 gm.
Ammonii Chloridi	.	.	gr. xij	.8 gm.
Aquæ	.	.	℥jss	45 cc.

Fiat haustus. Mitte vi tales. Solve.

Sig.—Bis, ter, quaterve in die sumendus.

180

R Liquoris Sodæ Chlorinatæ	℥iss	5.3 cc.
Syrupi Tolutani	℥j	30 cc.
Tinct. Serpentariæ	℥vj	21 cc.
Aquæ ad	℥viii	230 cc.

Fiat mistura. Misce.

Sig.—An eighth part every six hours.

181

R Vini Antimonialis	℥ss	1.8 cc.
Liquoris Ammonii Acetatis	℥iss	45 cc.
Magnesii Sulphatis	℥j	30 gm.
Aquæ Camphoræ ad	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat coch. mag. ter in die.

Relapsing Fever—

182

R Potassii Nitratis	.	.	gr. cxx	8 gm.
Acidi Nitrici Diluti	.	.	℥j	3.5 cc.
Tinct. Digitalis	.	.	℥ss	1.8 cc.
Aquæ Rosæ ad	.	.	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—One ounce every four hours.

Scarlet Fever—

183

R Potassii Chloratis	.	.	℥jss	6 gm.
Acidi Hydrochlorici	.	.	℥xl	2.4 cc.
Glycerini	.	.	℥j	30 cc.
Aquæ ad	.	.	℥xxx	568 cc.

Fiat gargarisma. Solve et misce; evolve Cl₂.

Sig.—Utendum coch. mag. ex cyatho aquæ.

184

R	Liquoris Sodæ Chlorinatæ	℥jv	15 cc.
	Aquæ ad	℥viij	230 cc.
	Fiat gargarisma. Misce.		
	Sig.—Utendum ℥ss ex aquâ si opus sit.		

Syphilis (constitutional), for Local Treatment see p. 98.

185

R	Hydrargyri cum Cretâ	gr. iij	.2 gm.
	Pulveris Opii	gr. 4	.016 gm.
	Fiat pilula. Mitte vii tales. Misce.		
	Sig.—Sumat unam nocte manequē.		

186

	Hydrargyri Perchloridi	gr. ij	.13 gm.
	Ammonii Chloridi	gr. x	.65 gm.
	Extracti Sarsæ Liquidi	℥iij	90 cc.
	Liq. Sarsæ Co. Conc. ad	℥xij	340 cc.
	Fiat mistura. Solve et misce.		
	Sig.—Sumat ℥ss ter in die.		

187

R	Hydrargyri Perchloridi	gr. j	.065 gm.
	Potassii Iodidi	gr. lx	4 gm.
	Tinct. Calumbæ	℥vi	21 cc.
	Aquæ ad	℥vj	170 cc.
	Fiat mistura. Solve et misce.		
	Sig.—Habenda ℥ss ter in die post cibum.		

188

R	Potassii Iodidi	℥vi	22 gm.
	Liquoris Hydrargyri Perchloridi	℥i	30 cc.
	Spiritūs Chloroformi	℥iii	10.5 cc.
	Aquæ ad	℥x	280 cc.
	Fiat mistura. Solve et misce.		
	Sig.—Capiat ℥iv ter in die ex aquâ post cibos.		

Infectious Diseases

189

R Potassii Iodidi gr. v .32 gm.
 Ammonii Carbonatis gr. vi .4 gm.
 Liquoris Hydrargyri Perchloridi ʒi 3.5 cc.
 Infusi Gentianæ Compositi ad ʒi 30 cc.
 Fiat haustus. Mitte viii tales. Solve et misce.
 Sig.—Sumat ter in die ex aquâ post cibos. P.P.A.
Syphilitic Stomatitis.

190

R Hydrargyri Perchloridi gr. ij .13 gm.
 Acidi Nitrici Diluti ℥xxx 1.8 cc.
 Tinct. Myrrhæ ʒj 30 cc.
 Aquæ ad ʒviij 2.30 cc.

Fiat garg. Misce.

Sig.—Utendum coch. mag. ex cyatho amplo
 aquæ t. i. d.

Syphilitic Stomatitis.

Tetanus—

191

R Magnesii Sulphatis ʒi 30 gm.
 Aquæ Destillatæ ʒiv 115 cc.

Fiat injectio. Solve.

Sig.—ʒss to be injected into the cerebro-spinal fluid
 by lumbar puncture every six hours.

192

R Acidi Carbolicî ℥x .6 cc.
 Aquæ Destillatæ ʒi 30 cc.

Fiat injectio hyp. Misce.

Sig.—ʒss to be injected every two hours
 necessary.

Typhoid Fever—

193

R Tinct. Catechu	-	-	℥ss	1.8 cc.
Tinct. Opii	-	-	℥x	.6 cc.
Olei Terebinthinæ	-	-	℥xv	.9 cc.
Mucilaginis Acaciæ	-	-	℥ij	7 cc.
Tinct. Chloroformi Compositæ	-	-	℥xx	1.2 cc.
Aquæ Menthæ Piperitæ ad	-	-	℥j	30 cc.
Fiat haustus. Mitte vi tales. Solve et misce.				
Sig.—Sumendus hora somni. P.P.A.				

In hæmorrhagæ.

194

R Olei Terebinthinæ	-	-	℥ij	7 cc.
Tinct. Lavandulæ Compositæ	-	-	℥ss	15 cc.
Sacchari Purificati	-	-	℥ij	8 gm.
Pulveris Acaciæ	-	-	℥ij	8 gm.
Aquæ ad	-	-	℥vj	170 cc.

Fiat mistura. Misce

Sig.—Capiatur ℥ss quartâquâque horâ si opus sit. P.P.A.

In hæmorrhagæ.

195

R Tinct. Camphoræ Compositæ	℥iij	10.5 cc.
Pulveris Cretæ Aromatici	℥ij	8 gm.
Syrupi Simplicis	℥ss	15 cc.
Aquæ Camphoræ ad	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ℥j post sedes liquidas. P.P.A.

In diarrhæa.

196

R Argenti Nitratis	-	-	gr. iij	.2 gm.
Pulveris Opii	-	-	gr. vj	.4 gm.
Pulveris Ipecacuanhæ	-	-	gr. vj	.4 gm.
Ung. Kaolin	-	-	q.s.	

Fiat massa. Divide in pilulas duodecem. Misce.

Sig.—One every four or six hours.

In diarrhæa.

INTOXICATIONS.

Alcoholism (acute)—

197

R Chloralis Hydratis	-	-	̄jss	6 gm.
Potassii Bromidi	-	-	̄ij	8 gm.
Spiritus Ætheris Co.	-	-	̄ij	7 cc.
Tinct. Valerianæ	-	-	̄iij	10.5 cc.
Aquæ ad	-	-	̄vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ̄ss secunda vel tertiâquâque horâ.

In threatened delirium.

198

R Liquoris Arsenicalis	-	-		
Tinct Capsici	-	-	āā ℥j	.06 cc.
Aquæ ad	-	-	̄j	30 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Take every hour after food.

In vomiting.

199

R Camphoræ Monobromidi	-	gr. v	.32 gm.
Confectionis Rosæ	-	q.s.	q.s.

Fiat pilula. Mitte xii tales. Misce.

Sig.—Sumat j vel ij si opus sit.

In sleeplessness.

200

R Potassii Bromidi	-	-	̄jss	6 gm.
Tinct. Digitalis	-	-	̄j	3.5 cc.
Tinct. Capsici	-	-	̄ss	1.8 cc.
Aquæ Camphoræ ad	-	-	̄vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ̄j ter in die.

In excitement.

Alcoholism (Chronic)—

201

R Atropinae Sulphatis	-	-	gr. $\frac{1}{2}$.016 gm.
Strychninae Hydrochloridi	-	-	gr. $\frac{1}{2}$.032 gm.
Pulveris Capsici	-	-	gr. x.	.65 gm.
Fiant pilulae xxiv. Misce.				

Sig.—Take one thrice a day for a week, then two thrice a day for another week with the following mixture.

202

R Tinct. Cinchonae Compositae	̄iv	15 cc.
Spiritus Ammoniae Aromatici	-	-
-	̄vi	21 cc.
Spiritus Chloroformi	-	-
-	̄iv	15 cc.
Aquae Camphorae	-	-
-	̄viii	230 cc.
Fiat mistura. Misce.		

Sig. Capiat ̄ss ex aqua t.i.d.
(M'Bride.)

Arsenic Poisoning—

203

R Liq. Ferri Perchlor. Fort.	̄i	3.5 cc.
Aquae ad	-	-
-	̄ij	60 cc.
Misce.		

Sig. No. 1.

R Magnesii Hydroxidi	-	-	̄i	4 gm.
Aquae ad	-	-	̄iiss	75 cc.

Solve

Sig. No. 2.—Mix with No. 1 and give ̄ss every 5 or 10 minutes until the symptoms are relieved.

Lead Poisoning (see also 25 and 26)

204

R Magnesii Sulphatis	-	-	5ij	8 gm.
Potassii Iodidi	-	-	gr. iij	.2 gm.
Aquæ ad	-	-	5i	30 cc.

Fiat haustus. Mitte viii tales. Solve.
 Sig.—Sumendus t. i. d. post cibos.

Morphia or Opium Poisoning—

205

R Sol. Potassii Permanganatis
 (1 in 1,000) - - - Oi 568 cc.
 Sig.—Use to wash out the stomach. Afterwards
 give the following pill.

R Liq. Atropinæ Sulphatis	-	℥ii	.12 cc.
Liq. Strychninæ Hydrochlor.	℥iv	.24 cc.	
Aquæ Destillatæ ad	-	5i	30 cc.

Fiat haustus. Mitte sex tales. Misce.
 Sig.—Capiat quartis horis donec æger
 recuperatus sit.

Strychnine Poisoning—

206

R Apomorphinæ Hydrochloridi	gr. i	.065 gm.
Acidi Hydrochlorici Diluti	℥i	.06 cc.
Aquæ Destillatæ ad	-	℥c 6 cc.

Fiat injectio hyp. Solve et misce.

Sig.—Five to ten drops to be used promptly after washing out the stomach with permanganate solution as above. Then give 5ii of bromide of potassium every hour if necessary.

NERVOUS DISEASES.

Chorea—

207

R Tincturæ Ferri Perchloridi	ʒii	7 cc.
Liquoris Arsenici Hydroch-		
lorici - - - -	ʒi	3.5 cc.
Glycerini - - - -	ʒvi	21 cc.
Aquæ ad - - - -	ʒvi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒii ter in die post cibos ex aquâ.

208

R Chloralis Hydratis - -	ʒi	4 gm.
Sodii Bromidi - - -	ʒij	8 gm.
Syrupi Pruni Virginianæ -	ʒvi	21 cc.
Aquæ Menthæ Piperitæ ad	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat cochleare plenum quartis vel sextis
horis more dicto.*In sleeplessness.*

Debility—

209

R Acidi Phosphorici Diluti -	ʒjss	5.3 cc.
Tinct. Nucis Vomicae -	ʒiss	5.3 cc.
Tinct. Cinchonæ Compositæ	ʒij	7 cc.
Infusi Aurantii ad - - -	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Coch. magnum ter in die sum.

210

R Quininæ Sulphatis	-	-	gr. xij	.8 gm.
Acidi Phosphorici Diluti	-	-	ʒjss	5.3 cc.
Syrupi Aurantii	-	-	ʒvj	21 cc.
Aquæ ad	-	-	ʒviij	230 cc.

Fiat mistura. Solve et misce.

Sig. Coch. magna duo ter in die sum.

Epilepsy—

211

R Ammonii Bromidi	-	-		
Sodii Bromidi	-	-		
Sodii Biboratis	-	-	āā ʒvi	22 gm.
Aquæ Camphoræ ad	-	-	ʒx	280 cc.

Fiat mistura. Solve.

Sig.—Capiat ʒiv ter in die post cibos.

212

R Sodii Bromidi	-	-		
Potassii Bromidi	-	-		
Ammonii Bromidi	-	-	āā ʒi	4 gm.
Syrupi Simplicis	-	-	ʒi	30 cc.
Aquæ Chloroformi ad	-	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. largum quartis horis post cibos.

213

R Potassii Iodidi	-	-	gr. xxiv	1.6 gm.
Potassii Bromidi	-	-	gr. xxx	2 gm.
Ferri et Ammonii Citratis	-	-	gr. xxx	2 gm.
Aquæ Anethi ad	-	-	ʒvj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumenda ʒj ter in die post cibos.

With anæmia.

214

R Potassii Bromidi	• •	3ij	8 gm.
Ferri et Ammonii Citratis	• •	3j	4 gm.
Spiritus Ammoniae Aromatici		3ij	7 cc.
Infusi Calumbae ad	• •	5vj	170 cc.

Fiat mistura. Solve et misce.

Sig. = Capienda 5j ter in die post cibos.

With anaemia.

Hiccough—

215

R Chloralis Hydratis	• •		
Potassii Bromidi	• •	5ā 3i	4 gm.
Aquae Menthae Piperitae ad		5vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat 5ss ter in die.

Hysteria—

216

R Spiritus Ammoniae Fetidi	• •	5i	30 cc.
Tincturae Valerianae Am-			
moniatæ	• •	5iss	45 cc.
Spiritus Ammoniae Aromatici	• •	5vi	21 cc.
Aquae Menthae Piperitae ad		5viii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat 5iv ter in die.

217

R Tinct. Asafetidae	• •		
Tinct. Sumbul	• •	5ā 5ij	7 cc.
Tinct. Opii	• •	5ss	1.8 cc.
Aquae Menthae Piperitae ad		5viii	230 cc.

Fiat mistura. Misce.

Sig. = An eighth part every four hours.

218

R Tinct. Valerianæ Ammoni-

atæ - - - - -	℥jv	15 cc.
Spiritus Chloroformi - - -	℥ij	7 cc.
Infusi Valerianæ ad - - -	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ℥j pro re nata.

Insomnia—

219

R Potassii Bromidi - - -	℥i	4 gm.
Chloralis Hydratis - - -	℥ss	2 gm.
Syrupi Aurantii - - -	℥vi	21 cc.
Aquæ ad - - -	℥iij	90 cc.

Fiat haustus. Misce.

Sig. Capiat dimidium horæ somni et residuum post horas quatuor, si opus sit.

220

R Paraldehydi - - -	℥iss	5.3 cc.
Syrupi Limonis - - -	℥ij	7 cc.
Misturæ Amygdalæ ad - -	℥i	30 cc.

Fiat haustus. Misce.

Sig. Sumendus horæ somni. P.P.A.

In cardiac cases.

221

R Sulphonalis - - -	gr. xv	1 gm.
Syrupi - - -	℥i	3.5 cc.
Pulveris Tragacanthæ Co. -	gr. x	.65 gm.
Aquæ Chloroformi ad - -	℥i	30 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Sumendus una hora ante decubitum. P.P.A.

222

R Veronalis - - -	gr. v	.32 gm.
-------------------	-------	---------

Fiat pulvis. Mitte sex tales.

Sig.—One at night, with caution in heart cases.

223

R Trionalis gr. x .65 gm.

Fiat pulvis. Mitte sex tales.

Sig.—One to two when not due to pain.

224

R Tinct. Opii ʒj 3.5 cc.

Potassii Bromidi ʒjss 6 gm.

Syrupi Rhœados ʒj 30 cc.

Aquæ ad ʒvj 170 cc.

Fiat mistura. Misce.

Sig.—Sumat ʒj horâ somni vel ter in die.

When due to pain.

225

R Extracti Cannabis Indicæ . gr. ½ .032 gm.

Fiat pilula sec. art. Mitte xii tales. Misce.

Sig.—Take one at bed-time.

When due to headache.

Nervous Excitability—

226

R Chloralis Hydratis

Potassii Bromidi āā gr. x .65 gm.

Syrupi Aurantii ʒss 15 cc.

Aquæ ad ʒj 30 cc.

Fiat haustus. Mitte vi tales. Solve et misce.

Sig.—To be taken at bed-time. May be given thrice daily.

Neuralgia—

227

R Tinct. Gelsemii ʒxv .9 cc.

Aquæ Camphoræ ʒj 30 cc.

Fiat haustus. Mitte vi tales. Misce.

Sig.—Sumat quârtâquâque horâ si opus sit.

Dental.

228

R Tinct. Gelsemii	-	-	℥vj	.35 cc.
Tinct. Ferri Perchloridi	-	-	℥x	.6 cc.
Syrupi Simplicis	-	-	℥ss	1.8 cc.
Aquæ ad	-	-	℥j	30 cc.
Fiat haustus. Mitte viii tales. Misce.				
Sig.—Repetatur ter in die post cibum.				

Dental.

229

R Phenacetini	-	-	gr. v	.32 gm.
Caffeinæ Citratis	-	-	gr. ii	.13 gm.
Quininæ Sulphatis	-	-	gr. iii	.2 gm.

Fiat pulvis. Mitte vi tales. Misce.

Sig.—Capiat unum tertiis horis dolore argenti.

Migraine.

230

R Phenacetini	-	-	gr. x	.65 gm.
Phenazoni	-	-	gr. v	.32 gm.
Caffeinæ Citratis	-	-	gr. iii	.2 gm.

Fiat pulvis. Mitte vi tales. Misce.

Sig.—Capiat unum quater in die.

Migraine.

231

R Ammonii Chloridi	-	-	℥iii	12 gm.
Tincturæ Gelsemii	-	-	℥i	3.5 cc.
Tincturæ Aurantii	-	-	℥ii	7 cc.
Aquæ Chloroformi ad	-	-	℥vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ℥iv dnabus horis.

Facial.

232

R Butyl-Chloralis Hydratis	-	gr. iij	.2 gm.
Gelsemininæ Hydrochloridi	-	gr. $\frac{1}{30}$.0022 gm.
Mucilaginis Acaciæ	-	q.s.	q.s.

Fiat pilula. Mitte xii tales. Misce

Sig.—Sumat unam tertiis horis donec dolor levatus sit.

Facial.

233

R Quininae Valerianatis -	-	gr. x	.65 gm.
Tinct. Sumbul -	-	ʒij	7 cc.
Extracti Taraxaci Liquidi -	-	ʒvj	21 cc.
Infusi Cascarillae ad -	-	ʒvj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ʒji ter in die.

Hysterical.

234

R Chloralis Hydratis -	-	ʒss	15 gm.
Camphoris -	-	ʒiij	12 gm.
Mentholis -	-	ʒi	4 gm.

Fiat pigmentum. Misce bene.

Sig.—Pinge ad dolentem partem ope penicilli.

235

R Amyl-Hydridi -	-	-	-
Alcoholis Absoluti -	-	āā ʒij	7 cc.
Aconitinæ -	-	gr. i	.065 gm.
Veratrinæ -	-	gr. v.	.32 gm.
Collodii ad -	-	ʒi	30 cc.

Fiat pigmentum. Misce.

Sig.—Applicandum affectis partibus ope penicilli
si necessitas sit

236

R Chloroformi -	-	ʒvj	21 cc.
Ætheris -	-	ʒj	30 cc.
Spiritus Camphoræ -	-	ʒiij	90 cc.
Tinct. Opii -	-	ʒjss	5.3 cc.

Fiat lotio. Misce.

Sig.—Apply to painful part on lint.

237

R Linimenti Belladonnæ -	-	-	-
Linimenti Aconiti -	-	-	-
Linimenti Chloroformi -	-	āā ʒi	30 cc.

Fiat linimentum. Misce.

Sig.—Infricandum leniter dolenti parti
si opus sit. P.P.A.

Neurasthenia—

238

R Quininæ Sulphatis	-	-	
Ferri Sulphatis	-	āā gr. xii	.8 gm.
Liq. Strychninæ Hydrochl.		℥xxx	1.8 cc.
Acidi Hydrobromici Diluti		ʒii	7 cc.
Aquæ Chloroformi ad	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat unciam bis in die post cibum.

239

R Liquoris Ferri Perchloridi	-	ʒii	7 cc.
Quininæ Sulphatis	-	ʒss	2 gm.
Liq. Strychninæ Hydrochl.		ʒi	3.5 cc.
Acidi Phosphorici Diluti	-	ʒii	7 cc.
Aquæ Chloroformi ad	-	ʒviii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat coch. mag. t.i.d. ex aquâ post cibos.

240

R Sodii Hypophosphitis	-	-	
Calcii do.	-	āā ʒi	4 gm.
Ferri do.	-	ʒss	2 gm.
Syrupi Simplicis	-	ʒij	60 cc.
Aquæ Chloroformi ad	-	ʒvi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat sextam partem ter in die post cibos.

Neuritis—

241

R Acetanilidi	-	ʒi	4 gm.
Caffeinæ	-	gr. xx.	1.3 gm.
Acidi Tartarici	-	gr. xxx	2 gm.
Sodii Bicarbonatis	-	gr. xlv.	3 gm.

Fiat pulvis. Misce.

Sig.—Capiat grana quinque vel decem dolore
urgenti.

242

R Mentholi	-	-	-	℥ss	2 gm.
Etheris	-	-	-	-	-
Chloroformi	-	-	-	āā ʒii	7 cc.

Fiat pigmentum. Misce.

Sig.—Applicandum affectis partibus more dicto.

Paralysis—

243

R Extracti Physostigmatis	-	gr. iii	.2 gm.
Pulveris Glycyrrhizæ	-	q.s.	-

Fiat massa. Divide in pilulas triginta. Misce.

Sig.—Capiat unam tertiis horis si opus sit.

244

R Strychninæ Sulphatis	-	gr. ii.	.13 gm
Aquæ Destillatæ	-	℥c	6 cc.

Fiat injectio hyp. Solve.

Sig.—Duo vel quatuor minima diebus alternis utenda.

Not in cerebral or spinal cases.

RESPIRATORY DISEASES.

Asthma —

245

R Tinct. Lobeliæ Ætherææ	-	ʒij	7 cc.
Spiritus Ætheris	-	ʒiij	10.5 cc.
Tinct. Conii	-	ʒij	7 cc.
Misturæ Amygdalæ ad	-	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ʒss ter in die. P.P.A.

246

R Tinct. Lobeliæ Æthereæ	-	̄iv	15 cc.
Tinct. Hyoscyami	-	̄vi	21 cc.
Spiritus Ætheris Co.	-	̄iv	15 cc.
Syrupi Tolutani ad	-	̄iv	115 cc.

Fiat mistura. Misce.

Sig.—Take one teaspoonful in water every half-hour during the paroxysm.

247

R Potassii Iodidi	-	-	̄ii	8 gm.
Spiritus Ammoniae Aromatici	̄iv			15 cc.
Tincturæ Lobeliæ Æthereæ	̄iii			10.5 cc.
Spiritus Chloroformi	-	-	̄iv	15 cc.
Aquæ ad	-	-	̄viii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat ̄iv quartis horis ex aquâ post cibum.

248

R Potassii Iodidi	-	-	̄ss	2 gm.
Spiritus Ammoniae Aromatici	̄ij			10.5 cc.
Tinct. Belladonnae	-	-	̄jss	5.3 cc.
Tinct. Cinchonae Compositae	̄vj			21 cc.
Aquæ Menthae Piperitæ ad	̄vj			170 cc.

Fiat mistura. Misce.

Sig.—Habeat ̄j ter in die post cibos ex aquâ.

249

R Pulveris Lobeliæ	-	-		
Pulveris Stramonii	-	-		
Pulveris Theae Nigræ	āā	̄ss		15 gm.
Potassii Nitratis	-	-		
Pulveris Anisi	-	-		
Pulveris Fœniculi	-	āā	̄j	30 gm.

Fiat pulvis. Misce.

Sig.—A small teaspoonful to be burned on a plate and the smoke inhaled.

250

R Ætheris	•	•	•	•	5j	30 cc.
Olei Terebinthinæ	•	•	•	•	3iii	10.5 cc.
Tincturæ Benzoini Co. ad	•	•	•	•	5ij	60 cc.

Fiat inhalatio. Misce.

Sig.—A teaspoonful in a pint of hot water and the steam inhaled.

251

R Folii Belladonnæ	•	•	•	•	gr. vj	.4 gm.
Folii Hyoscyami	•	•	•	•	•	•
Folii Stramonii	•	•	•	•	āā gr. iij	.2 gm.
Olei Eucalypti	•	•	•	•	℥½	.03 cc.
Extracti Opii	•	•	•	•	gr. ½	.016 gm.
Aquæ Laurocerasi	•	•	•	•	q.s.	•

Moisten the powdered leaves with the extract of opium dissolved in the cherry laurel water, then dry and make into a cigarette. Send six.

Bronchial Asthma—

252

R Potassii Iodidi	•	•	•	•	5ij	8 gm.
Ammonii Carbonatis	•	•	•	•	3j	4 gm.
Tinct. Lobeliæ Ætherææ	•	•	•	•	5ij	7 cc.
Spiritus Chloroformi	•	•	•	•	5iv	15 cc.
Vini Ipecacuanhæ	•	•	•	•	3j	3.5 cc.
Infusi Senegæ ad	•	•	•	•	5vj	170 cc.

Fiat mistura. Misce.

Sig.—A table-spoonful in a wine-glassful of water every four hours.

253

R Tinct. Camphoræ Compositæ	•	•	•	•	•	•
Spiritus Ætheris	•	•	•	•	āā 3iij	10.5 cc.
Tinct. Scillæ	•	•	•	•	3j	3.5 cc.
Syrupi	•	•	•	•	5ss	15 cc.
Aquæ ad	•	•	•	•	5vj	170 cc.

Fiat mistura. Misce.

Sig.—Capiat 5j ter in die.

254

R Oxy mellis Scillæ	-	-	℥i	3.5 cc.
Syrupi Papaveris	-	-	℥j	3.5 cc.
Tinct. Camphoræ Compositæ	℥ss			1.8 cc.
Tinct. Lobeliæ Æthereæ	-	-	℥xv	.9 cc.
Misturæ Ammoniaci ad	-	-	℥j	30 cc.
Fiat haustus. Mitte vi tales. Misce.				
Sig.—Capiendus dosis ter in die.				

Bronchitis (acute)—

255

R Potassii Bicarbonatis	-	-	℥ii	8 gm.
Vini Ipecacuanhæ	-	-	℥iiss	8.8 cc.
Liquoris Ammonii Acetatis				
Syrupi Tolutani	-	-	℥i	30 cc.
Aquæ Camphoræ ad	-	-	℥vi	170 cc.
Fiat mistura. Solve et misce.				
Sig.—Capiat ℥iv tertiis horis.				

256

R Liquoris Ammonii Acetatis	℥vj			21 cc.
Spiritus Ammonizæ Aromat.	℥iij			10.5 cc.
Syrupi Aurantii	-	-	℥j	30 cc.
Aquæ ad	-	-	℥vj	170 cc.
Fiat mistura. Misce.				
Sig.—Sumat ℥j ter in die.				

257

R Liquoris Ammonii Acetatis	℥ss			15 cc.
Vini Ipecacuanhæ	-	-	℥j	3.5 cc.
Potassii Nitratis	-	-	℥ss	2 gm.
Syrupi	-	-	℥j	30 cc.
Sacchari Usti	-	-	℥ss	2 gm.
Aquæ Camphoræ ad	-	-	℥vj	170 cc.
Fiat mistura. Solve et misce.				
Sig.—Capiat ℥j ter in die.				

Bronchitis (chronic).

258

R Ammonii Carbonatis	-	-	3ss	2 gm.
Spiritus Ætheris	-	-	3ij	10.5 cc.
Tinct. Scillæ	-	-	3jss	5-3 cc.
Tinct. Camphoræ	Com-			
positæ	-	-	3ij	10.5 cc.
Tinct. Lavandulæ	Com-			
positæ	-	-	3jv	15 cc.
Infusi Senegæ ad	-	-	5viij	230 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat 5j ter in die.

With scanty expectoration.

259

R Ammonii Carbonatis	-	-	3i	4 gm.
Tincturæ Nucis Vomicae	-	-	3iss	5.3 cc.
Spiritus Chloroformi	-	-	3iii	10.5 cc.
Infusi Senegæ ad	-	-	5vi	170 cc.

Fiat mistura. Solve et misce.

Sig. Capiat 3iv tertiis horis.

In debility.

260

R Ammonii Carbonatis	-	-	3ss	2 gm.
Spiritus Etheris	-	-	3ij	7 cc.
Syrupi	-	-	5j	30 cc.
Infusi Gentianæ Co. ad	-	-	5vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat 5j quartâquâque horâ.

In debility.

261

R Ammonii Carbonatis	-	-	3j	4 gm.
Tinct. Digitalis	-	-	3j	3.5 cc.
Tinct. Camphoræ	Com-			
positæ	-	-	3ij	10.5 cc.
Misturæ Amygdalæ ad	-	-	5vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat 5j ter in die. P.P.A.

With weak heart.

262

R Tinct. Guaiaci Ammoniatae	℥iij	10.5 cc.
Mucilaginis Tragacanthæ	℥iij	90 cc.
Infusi Senegæ ad	℥viiij	230 cc.

Fiat mistura. Misce.

Sig.—Coch. mag. duo vel tria ter in die capienda.
P.P.A.*In rheumatic cases.*

263

R Tinct. Scillæ	-	-	-
Acidi Sulphurici Diluti	āā	℥jss	5.3 cc.
Tinct. Opii	-	℥ss	1.8 cc.
Infusi Cascarillæ ad	-	℥vj	170 cc.

Fiat mistura. Misce.

Sig. Sumenda ℥ss ter in die.

With excessive secretion.

264

R Acidi Tannici	-	℥ss	2 gm.
Acidi Nitrici Diluti	-	℥j	3.5 cc.
Tinct. Camphoræ Co.	-	℥ij	7 cc.
Infusi Gentianæ Co. ad	-	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capienda ℥j ter in die.

As above.

265

R Ammonii Chloridi	-	gr. x	.65 gm.
Spiritus Chloroformi	-	℥xx	1.2 cc.
Vini Antimonialis	-	℥x	.6 cc.
Aquæ Camphoræ ad	-	℥j	.30 cc.

Fiat haustus. Mitte vi tales. Solve et misce.

Sig. Sumat ter in die.

With rapid pulse.

266

R Spiritus Ætheris	-	-	5jss	5.3 cc.
Spiritus Ammoniae Aromatici	5ij			7 cc.
Infusi Aurantii ad	-	-	5vj	170 cc.

Fiat mistura. Misce.

Sig.—Capiat 5j ter in die.

With difficult breathing.

267

R Ammonii Chloridi	-	-		
Ammonii Carbonatis	āā	gr. xv		1 gm.
Pulveris Ipecacuanhæ	-	gr. iij		.2 gm.
Morphinæ Hydrochloridi	-	gr. j		.065
Mucilaginis Acaciæ	-	q.s.		q.s.

Fiat massa. Divide in pilulas decem. Misce.

Sig.—Sumat unam omni nocte maneque.

To assist expectoration.

268

R Pilulæ Scillæ Compositæ	gr. v			.32 gm.
---------------------------	-------	--	--	---------

Mitte xii tales.

Sig.—Capiat unam nocte maneque.

To assist expectoration.

Bronchiectasis—

269

R Iodoformi	-	-	gr. xx.	1.5 gm.
Olei Eucalypti	-	-	5ss	15 cc.
Spiritus Rectificati ad	-	-	5i	30 cc.

Fiant guttæ. Solve et misce.

Sig.—A few drops on the sponge of a dry inhaler.

270

R Olei Eucalypti	-	-	3ss	1.8 cc.
Olei Terebinthinæ	-	-	5iv	15 cc.
Magnesii Carbonatis Levis	5iij			12 gm.
Aquæ ad	-	-	5x	280 cc.

Fiat inhalatio. Misce.

Sig.—Coch mag. duo ex aquæ calidæ octario.

Catarrh—

271

R	Liquoris Ammonii Acetatis	℥j	30 cc.
	Ammonii Carbonatis	- - 3ss	2 gm.
	Potassii Nitratis	- - 3j	4 gm.
	Aquæ Camphoræ ad	- - 5vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat 5j ter in die.

When feverish.

272

R	Pulveris Opii	- - - gr. j	.065 gm.
	Potassii Nitratis	- - - gr. viij	.52 gm.
	Pulveris Ipecacuanhæ	- - - gr. j	.065 gm.

Fiat pulvis. Mitte iii tales. Misce.

Sig.—Unus hora somni sumendus.

In early stage.

273

R	Cocainæ Hydrochloridi	- - - gr. jss	.1 gm.
	Mentholis	- - - gr. ijss	.16 gm.
	Acidi Salicylici	- - - gr. viij	.52 gm.
	Acidi Borici	- - - 3j	4 gm.
	Pulveris Amyli	- - - 5jss	6 gm.

Fiat insufflatio. Misce.

Sig.—A pinch every two hours as a snuff
after blowing the nose.*In nasal type.*

Cough—

274

R	Liq. Morphine Hydrochlor.	3i	3.5 cc.
	Chloroformi	- - - 3ss	1.8 cc.
	Tinct. Cannabis Indicæ	- 3i	3.5 cc.
	Pulveris Tragacanthæ Co.	- 3i	4 gm.
	Spiritus Ætheris	- - 3iii	10.5 cc.
	Acidi Hydrocyanici Diluti	℥ xii	.7 cc.
	Aquæ ad	- - - 5vi	170 cc.

Fiat mistura. Misce.

Sig.—Uncia tussi urgente sumenda si opus sit.

P.P.A.

In irritable cough.

275

R Morphinae Hydrochloridi	-	gr. v	.32 gm.
Atropinae Sulphatis	-	gr. $\frac{1}{16}$.0065 gm.
Glycerini ad	-	$\bar{\text{ss}}$ i	30 cc.
Solve et adde.			
Chloroformi	-	$\bar{\text{ss}}$ ss	5.3 cc.
Olei Menthae Piperitae	-	\mathfrak{M} ij	.12 cc.
Alcoholis (90%)	-	q.s.	
Misce et adde.			
Alcoholis (90%) ad	-	$\bar{\text{ss}}$ ss	45 cc.
Fiant guttae. Misce.			
Sig.—Capiat v-xv guttas ex aqua vel in sacchari frustillo. P.P.A.			
<i>As above.</i>			

276

R Acidi Hydrobromici Diluti	\mathfrak{M} xl	2.4 cc.
Liquoris Morphinae Acetatis	\mathfrak{M} xxiv	1.4 cc.
Acidi Hydrocyanici Diluti	\mathfrak{M} xii	.7 cc.
Vini Ipecacuanæ - - -	$\bar{\text{ss}}$ i	3.5 cc.
Syrupi Papav. Rubri - - -	$\bar{\text{ss}}$ i	30 cc.
Glycerini - - - - -	$\bar{\text{ss}}$ ii	60 cc.
Aquae Chloroformi ad - -	$\bar{\text{ss}}$ vi	170 cc.
Fiat mistura. Misce.		
Sig. — Cap. coch. mag. tussi urgente.		P.P.A.
<i>As above.</i>		

277

R Tinct. Belladonnae	-	$\bar{\text{ss}}$ j	3.5 cc.
Tinct. Conii	-	$\bar{\text{ss}}$ ij	7 cc.
Vini Ipecacuanhae	-	$\bar{\text{ss}}$ jss	5.3 cc.
Aquae Chloroformi ad	-	$\bar{\text{ss}}$ vj	170 cc.
Fiat mistura. Misce.			
Sig.—Sumat $\bar{\text{ss}}$ j ter in die.			
<i>With much expectoration.</i>			

278

R Tinct. Camphoræ Com-		
positæ - - - -	ʒiij	10.5 cc.
Vini Ipecacuanhæ - -	ʒj	3.5 cc.
Acidi Hydrocyanici Diluti	℥xij	.7 cc.
Syrupi Scillæ - - -	ʒss	15 cc.
Aquæ ad - - - -	ʒvj	170 cc.

Fiat mistura. Misce.

Sig.—Capiat ʒj ter in die. P.P.A.

In dry cough.

279

R Codeinæ - - - -	gr. ss	.032 gm.
Pilulæ Galbani Compositæ	gr. v	.32 gm.

Fiat pilula. Mitte vi tales. Misce.

Sig.—One to be taken at bed-time.

*In night cough.*Hæmoptysis (*see also* 295 to 296)—

280

R Olei Terebinthinæ - -	℥xv	.9 cc.
Misturæ Amygdalæ ad -	ʒj	30 cc.

Fiat haustus. Mitte viii tales. Misce.

Sig.—To be taken every hour. P.P.A.

Laryngitis (*see also* 300 to 302)—

281

R Tincturæ Benzoini Com-		
positæ - - - -	ʒi	30 cc.

Fiat inhalatio. Misce.

Sig.—A teaspoonful to be used in a pint of hot water with an inhaler.

282

R Creosoti	· · · ·	̄iiss	5.3 cc.
Magnesii Carbonatis Levis		̄i	4 gm.
Aquæ ad	· · · ·	̄iij	60 cc.
Fiat inhalatio. Misce.			

Or

238

R Olei Pini Sylvestris	· · ·	̄i	3.5 cc.
Magnesii Carbonatis Levis		̄i	4 gm.
Aquæ ad	· · · ·	̄iij	60 cc.
Fiat inhalatio. Misce.			

Or

284

R Olei Eucalypti	· · ·	̄iij	7 cc.
Olei Pini Sylvestris	· · ·	̄iiss	5.3 cc.
Magnesii Carbonatis Levis		̄i	4 gm.
Aquæ ad	· · · ·	̄iij	60 cc.
Fiat inhalatio. Misce.			

Sig.—One teaspoonful to be put in a pint of boiling water and the steam inhaled. P.P.A.

Œdema of the Lung—

285

R Antimonii Tartarati	· · ·	gr. j	.065 gm.
Potassii Nitratis	· · ·	̄iij	3 gm.
Tinct. Hyoscyami	· · ·	̄iiss	5.3 cc.
Tinct. Digitalis	· · ·	̄iiss	1.8 cc.
Misturæ Amygdalæ ad	· · ·	̄viij	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ̄ij omnibus horis. P.P.A.

Phthisis—

286

R Codeinæ Sulphatis	-	-	gr. iij	.2 gm.
Liquoris Atropinæ Sulphatis	℥xiij			.7 cc.
Syrupi Tolutani	-	-	℥jss	45 cc.
Infusi Rosæ Acidi ad	-	-	℥vj	170 cc.
Fiat mistura. Solve et misce.				

Sig.—Sumat ℥ss quartâquâque horâ ex aquâ
tussi urgente.

Cough sedative.

287

R Liq. Morphinæ Hydroch-				
loridi	-	-	℥jss	5.3 cc.
Acidi Phosphorici Diluti	-	-	℥j	3.5 cc.
Acidi Hydrocyanici Diluti	-	-	℥xiij	.7 cc.
Syrupi Rhoëados	-	-	℥ss	15 cc.
Aquæ ad	-	-	℥jss	45 cc.
Fiat mistura. Misce.				

Sig.—Sumat ℥j tussi urgente ex aquâ. P.P.A.

Cough sedative.

288

R Calcii Hypophosphitis	-	-	℥j	4 gm.
Tinct. Camphoræ Co.	-	-	℥iij	10.5 cc.
Acidi Hydrocyanici Diluti	-	-	℥xiij	.7 cc.
Syrupi Simplicis	-	-	℥j	30 cc.
Infusi Columbæ ad	-	-	℥vj	170 cc.
Fiat mistura. Solve et misce.				

Sig.—Capiat ℥j ter in die. P.P.A.

As a tonic.

289

R Quininae Sulphatis	-	-	
Ferri Sulphatis	-	āā gr. viii	.52 gm.
Liq. Strychninae Hydrochlor.	℥xxiv		1.4 cc.
Acidi Sulphurici Diluti	-	℥xl	2.4 cc.
Syrupi Simplicis	-	ḡi	30 cc.
Infusi Calumbæ ad	-	ḡviii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat cochlearia magna duo t. i. d.
post cibos.

As a tonic.

290

R Quininae Sulphatis	-	-	
Pulveris Digitalis	-	āā gr. xij	.8 gm.
Pulveris Opii	-	- gr. vj	.4 gm.
Fiat pulvis.	Divide in duodecim chartas.	Misce.	

Sig.—Give one every four hours.

(Niemeyer's powders.)

With high temperature.

291

R Zinci Oxidi	-	-	gr. ii	.13 gm.
Extracti Belladonnae Viridis	-	-	gr. ¼	.016 gm.
Extracti Hyoscyami	-	-	gr. ¼	.032 gm.

Fiat pilula. Mitte xii tales. Misce.

Sig.—Capiat unam vel duas horā somni.

In night sweats.

292

R Acidi Gallici	-	-	gr. vj	.4 gm.
Extracti Cannabis Indicae	-	-	gr. ss	.032 gm.
Confectionis Rosae	-	-	q.s.	

Fiat pilula. Mitte xii tales. Misce.

Sig.—One to be taken every night.

In night sweats.

293

R Syrupi Papaveris	-	-	℥jv	15 cc.
Bismuthi Subnitratis	-	-	℥j	4 gm.
Mucilaginis Tragacanthæ	-	-	℥i	30 cc.
Aquæ ad	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ℥j ter in die. P.P.A.

In diarrhœa.

294

R Tincturæ Opii	-	-	℥i	3.5 cc.
Tincturæ Catechu	-	-	℥vi	21 cc.
Decocti Hæmatoxyli ad	-	-	℥vi	170 cc.

Fiat mistura. Misce.

Sig. Capiat coch. mag. ter in die
si opus sit.*In diarrhœa.*

295

R Magnesii Sulphatis	-	-	℥j	30 gm.
Acidi Gallici	-	-	℥j	4 gm.
Tinct. Digitalis	-	-	℥j	3.5 cc.
Extracti Ergotæ Liquidi	-	-	℥ij	7 cc.
Infusi Rosæ Acidi ad	-	-	℥vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Take one ounce every two or three hours.

In hæmoptysis.

296

R Extracti Ergotæ	-	-	gr. x	.65 gm.
Extracti Krameriæ	-	-	℥j	4 gm.
Extracti Belladonnæ	-	-	gr. v	.32 gm.
Pulveris Digitalis	-	-	gr. x	.65 gm.

Fiat massa. Divide in pilulas viginti. Misce.

Sig.—Capiat duas vel tres si opus sit.

In hæmoptysis.

297

R Creosoti	-	-	-	℥vij	.4 cc.
Iodoformi	-	-	-	gr. vj	.4 gm.
Acidi Carbolici (cryst.)	-	-	-	gr. viij	.52 gm.
Olei Theobromatis	-	-	-	q.s.	q.s.

Fiat massa. Divide in triginta pilulas. Misce.

Sig.—One pill every three hours.

In offensive sputum.

298

R Glycerini Acidi Carbolici	-	ʒiij	10.5 cc.
Chloroformi	-	℥viij	.5 cc.
Succi Conii	-	ʒii	7 cc.
Aquæ ad	-	ʒviij	230 cc.

Fiat inhalatio. Misce.

Sig.—An ounce in hot water.

In offensive sputum.

299

R Creosoti	-	-	-	-	-
Acidi Carbolici	-	-	-	-	-
Eucalyptolis	-	-	ʒā ʒii	7 cc.	
Spiritus Ætheris	-	-	-	-	-
Tinct. Iodi	-	-	ʒā ʒi	3.5 cc.	
Glycerini ad	-	-	ʒi	30 cc.	

Fiant guttæ. Misce.

Sig.—To be used continuously with a dry inhaler.

(Veo's.)

300

R Creosoti (Beechwood)-	-	ʒi	3.5 cc.
Olei Cassiæ -	-	℥xx	1.2 cc.
Eucalyptolis	-	ʒss	1.8 cc.
Paraffini Liquidil ad	-	ʒii	60 cc.

Fiat nebula. Misce.

Sig.—To be used with an atomiser.

In laryngeal cases.

301

R Iodi	-	-	-	gr. viii	.52 gm.
Mentholis	-	-	-	gr. x	.65 gm.
Creosoti (Beechwood)	-	-	-	℥xv	.9 cc.
Eucalyptolis	-	-	-	℥xx	1.2 cc.
Olei Cassiæ	-	-	-	℥v	.3 cc.
Paraffini Liquidī ad	-	-	-	̄i	30 cc.

Fiat nebula. Solve et misce.

Sig.—To be used with an atomiser.

In laryngeal cases.

302

R Cyllini	-	-	-	̄i	30 cc.
-----------	---	---	---	----	--------

Sig.—A few drops to be used with a dry inhaler.

In laryngeal cases.

303

R Cocainæ Hydrochloridi	-	gr. x	.65 gm.
Acid Borici	-	gr. jv	.26 gm.
Glycerini	-	℥xv	.9 cc.
Aquæ Destillatæ ad	-	̄j	30 cc.

Fiat applicatio. Solve et misce.

Sig.—Use with a throat brush or atomiser.

In dysphagia.

Pleurisy—

304

R Tinct. Veratri Viridis	-	̄iv	15 cc.
--------------------------	---	-----	--------

Sig.—Five drops to be taken every three hours increasing the tincture by a drop till the pulse is sufficiently lowered.

In early stage.

305

R Iodi	-	-	̄i	4 gm.
Potassii Iodidi	-	-	̄ss	2 gm.
Glycerini	-	-	̄ij	7 cc.
Spiritus Rectificati ad	-	-	̄j	30 cc.

Fiat pigmentum. Misce.

Sig.—Apply with brush nightly until the skin gets tender.

In chronic form.

Pneumonia—

306

R Potassii Nitratis	-	-	̄ij	4 gm.
Liquoris Ammonii Acetatis	-	-	̄jss	45 cc.
Spiritus Ætheris Nitrosi	-	-	̄iv	15 cc.
Tincturæ Aurantii	-	-	̄iii	10.5 cc.
Aquæ Camphoræ ad	-	-	̄vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ̄iv tertiis horis.

In febrile stage.

307

R Potassii Nitratis	-	-	̄j	4 gm.
Liq. Ammonii Acetatis	-	-	̄jss	45 cc.
Vini Ipecacuanhæ	-	-	̄j	3.5 cc.
Spiritus Ammonia Aromatici	-	-	̄ij	7 cc.
Decocti Hordei ad	-	-	̄vj	170 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ̄j ter in die.

In febrile stage.

308

R Tinctura Veratri Viridis	-	-	℥xl	2.4 cc.
Spiritus Etheris Nitrosi	-	-	̄vi	21 cc.
Potassii Citratis	-	-	̄ij	8 gm.
Syrupi Zingiberis	-	-	̄ss	15 cc.
Aquæ ad	-	-	̄vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare plenum tertiis horis.

In febrile stage.

309

R Ammonii Carbonatis	-	-	gr. v	.32 gm.
Spiritus Chloroformi	-	-	℥x	.6 cc.
Vini Colchici	-	-	℥v	.3 cc.
Liquoris Ammonii Acetatis	-	-	̄ij	10.5 cc.
Aquæ Mentha Piperitæ ad	-	-	̄j	30 cc.

Fiat haustus. Mite vi tales. Solve et misce.

Sig.—To be taken every four hours. P.P.A.

In gouty cases.

310

R. Ammonii Carbonatis	-	-	gr. xxx	2 gm.
Tinct. Aconiti	-	-	℥xx	1.2 cc.
Tinct. Cinchonæ	-	-	℥vj	21 cc.
Aquæ Menthræ Piperitæ ad	-	-	℥vj	170 cc.
Fiat mistura. Solve et misce.				
Sig.—Habeat ℥j ter in die.				

In later stages.

311

R. Quininae Sulphatis	-	-	℥ss	2 gm.
Acidi Hydrobromici Diluti	-	-	℥iii	10.5 cc.
Caffeinae Citratis	-	-	℥i	4 gm.
Liq. Strychninae Hydroch-				
loridi	-	-	℥ss	1.8 cc.
Aquæ ad	-	-	℥vi	170 cc.
Fiat mistura. Solve et misce.				
Sig.—Capiat ℥iv ter in die post cibos.				

In convalescence.

SKIN DEASES.

Abscess (tuberculous).

312

R. Iodoformi	-	-	gr. x	.65 gm.
Etheris Sulphurici	-	-	℥iss	5.3 cc.
Paraffini Liquidi ad	-	-	℥ss	1.8 cc.
Fiat injectio. Solve et misce.				
Sig. To be injected after emptying the abscess.				

Acne Vulgaris—

313

R Pulveris Cretæ	-	-	-	gr. v	.32 gm.
β Naphtholis	-	-	-		
Camphoræ	-	-	-	āā gr. x	.65 gm.
Vaselini	-	-	-	ʒiiss	6 gm.
Saponis Viridis	-	-	-	gr. xv.	1 gm.
Sulphuris Præcipitati	-	-	-	gr. l	3.3 gm.
Fiat unguentum. Misce.					
Sig.—Utendum mane nocteque.					

314

R Thymolis	-	-	-	gr. x	.65 gm.
Sulphuris Præcipitati	-	-	-	ʒss	2 gm.
Paraffini Mollis ad	-	-	-	ʒi	30 gm.
Fiat ung. Misce.					
Sig.—Applicandum mane nocteque si opus sit.					

315

R Unguenti Thymolis Com-	-	-	-	ʒiiss	45 gm.
positi, B.P.C.	-	-	-	ʒss	15 gm.
Paraffini Mollis	-	-	-	ʒss	
Fiat unguentum. Misce.					
Sig.—Applicetur mane nocteque.					

Acne Rosacea—

316

R Calamina Præparata	-	-	-		
Cretæ Præparata	-	-	-	āā ʒi	4 gm.
Glycerini	-	-	-	ʒiij	10.5 cc.
Acidi Hydrocyanici Diluti	-	-	-	ʒss	1.8 cc.
Liquoris Calcis	-	-	-	ʒiij	90 cc.
Aquæ Rosæ ad	-	-	-	ʒviij	230 cc.
Fiat lotio. Misce.					
Sig.—To be dabbed on and allowed to dry. P.P (Sartin).					

317

R Sulphuris Præcipitati	-	℥ss	6 gm.
Glycerini	-	℥j	3.5 cc.
Spiritus Rectificati	-	℥ss	15 cc.
Aquæ Rosæ ad	-	℥jv	115 cc.

Fiat lotio. Misce.

Sig.—After steaming face, paint lotion on
at night. P.P.A.

318

R Glycerini Acidi Carbolici	-	℥ij	60 cc.
Sulphuris Præcipitati	-	℥ss	45 gm.
Spiritus Rectificati	-	℥ij	60 cc.

- Misce.

Sig.—To be applied every night. P.P.A.
(Hebra's Prescription).

Alopecia—

319

R Spiritus Rectificati	-	℥i	30 cc.
Tinct. Cantharidis	-	-	-
Tinct. Capsici	-	āā ℥ij	7 cc.
Spiritus Rosmarini	-	℥ij	60 cc.
Liquoris Ammoniz Fortis	-	℥ss	15 cc.

Fiat lotio. Misce.

Sig.—To be gently rubbed on the affected parts.

320

R Acidi Lactici	-	℥ss	15 cc.
Aquæ ad	-	℥iij	90 cc.

Fiat lotio. Misce.

Sig.—Infrietur leniter affectis partibus more dicto.

321

℞ Olei Amygdalæ Dulcis	-	
Liquoris Ammoniaë Fortis	āā	℥ss 15 cc.
Spiritus Rosmarini	-	℥ij 60 cc.
Aquæ Mellis ad	-	℥iv 115 cc.

Fiat lotio. Misce.

Sig.—Utenda nocte manequē.
(Erasmus Wilson's Lotion).

322

R Resorcini	-	℥jss 6 gm.
Olei Ricini	-	℥jss 45 cc.
Balsami Peruviani	-	gr. viij .52 gm.
Spiritus Rectificati ad	-	℥vi 170 cc.

Fiat lotio. Solve et misce.

Sig.—Applicetur affectis partibus nocte.

323

℞ Olei Sinapis Volatilis	-	℥j 3.5 cc.
Olei Ricini	-	℥ij 7 cc.
Spiritus Rosmarini ad	-	℥jv 115 cc.

Fiat lotio. Misce.

Sig.—Paint on the patches once or twice a day.

324

℞ Tinct. Cantharidis	-	℥j 3.5 cc.
Glycerini	-	-
Acidi Acetici Diluti	āā	℥jss 45 cc.
Spiritus Rosmarini	-	℥ij 60 cc.
Aquæ Rosæ ad	-	℥viiij 230 cc.

Fiat lotio. Misce.

Sig.—Applicanda omni nocte.

Boils—

325

℞ Pilulæ Calcis Sulphuratæ	-	gr. i .065 gm.
Mitte viginti tales.		

Sig.—Unam t.d.s. Sensim augenda dosis ad
octo per diem.

326

R Unguenti Resinæ	-	-	̄i	30 gm.
Unguenti Hydrargyri				
Aminon.	-	-	̄ss	15 gm.
Paraffini Mollis ad	-	-	̄ii	60 gm.

Fiat unguentum. Misce.

Sig.—Applicandum super linteam carptum extensum omni nocte.

Burns—

327

R Olei Eucalypti	-	-	̄ss	1.8 cc.
Olei Lini	-	-	̄iij	90 cc.
Liquoris Calcis	-	-	̄vi	170 cc.

Fiat lotio. Misce.

Sig.—To be applied on smooth linen and covered with cotton wool.
(Carron Oil).

328

R Acidi Picrici	-	-	̄ss	2 gm.
Aquæ ad	-	-	̄viii	230 cc

Fiat lotio. Solve.

Sig.—To be applied with strips of sterile gauze. After forty-eight hours remove, and wash with permanganate of potassium solution.

Cancer—

329

R Zinci Chloridi	-	-	̄i	30 gm.
Farinæ Tritici	-	̄ii vel	q.s	8 gm.
Liquoris Opii Sedativi	-	̄i		30 cc.

Fiat pasta sec. art. Misce.

Sig.—Applicandum more dicto.

330

R Coley's Fluid	• • •	℥x	.6 cc.
Aquæ Destillatæ	• • •	℥ss	15 cc.

Fiat inject. hyp. Misce.

Sig. — Minima sex quotidie vel diebus alternis
utenda, si opus sit.

In sarcoma.

Corns—

331

R Acidi Salicylici	• • •	ʒi	4 gm.
Extracti Cannabis Indicæ	• • •	gr. xv	1 gm.
Collodii (flexilis) ad	• • •	ʒi	30 cc.

Fiat pigmentum. Solve et misce.

Sig. Applicandum omni nocte, si opus sit.

Eczema

332

R Bismuthi Subnitratis	• • •	ʒjv	15 gm.
Zinci Oxidi	• • •	ʒj	4 gm.
Acidi Carbolici Liquefacti	• • •	℥ss	1.8 cc.
Vaselini Albi ad	• • •	ʒij	60 gm.

Fiat unguentum. Misce.

Sig. Utendum nocte maneque.

333

R Zinci Oxidi	• • •	ʒii	8 gm.
Acidi Salicylici	• • •	℥ss	2 gm.
Bismuthi Carbonatis	• • •	ʒi	4 gm.
Pulveris Amyli	• • •	ʒvi	22 gm.

Fiat pulvis. Misce.

Sig. To be dusted over the affected parts.

334

R Liqnoris Plumbi Subacetatis	℥j	3.5 cc.
Glycerini	- - - - - ℥ss	15 cc.
Aquæ ad	- - - - - ℥vj	170 cc.

Fiat lotio. Misce.

Sig. Applicanda affectis partibus.

335

R Liq. Carbonis Detergentis	- - - - - ℥ss	15 cc.
Acidi Hydrocyanici Diluti	℥j	3.5 cc.
Glycerini	- - - - - ℥ij	7 cc.
Aquæ ad	- - - - - ℥x	260 cc.

Fiat lotio. Misce.

Sig. - Applicanda si opus sit. P. P. A.

In painful forms.

336

R Plumbi Acetatis	- - - - - gr. x	.65 gm.
Zinci Oxidi	- - - - -	
Hydrargyri Subchloridi	āā gr. xx	1.3 gm.
Unguenti Hydrargyri Nitratis	gr. xxx	2 gm.
Adipis Recentis	- - - - - ℥iv	15 gm.
Olei Palmæ Purificati	- - - - - ℥ss	15 cc.

Fiat ung. Solve et misce.

Sig. Applicandum omni nocte.

When on the head.

337

R Boracis	- - - - - ℥j	4 gm.
Liquoris Plumbi Subacetatis	℥lxxx	1.8 cc.
Glycerini	- - - - - ℥ij	7 cc.
Aquæ ad	- - - - - ℥viij	230 cc.

Fiat lotio. Solve et misce.

Sig. Utenda sæpe.

When on the face.

338

R Zinci Sulphatis	-	-	gr. xxx	2 gm.
Zinci Oxidi	-	-	gr. lx	4 gm.
Pulveris Acaciæ	-	-	ʒij	8 gm.
Aquæ Sambuci ad	-	-	ʒxxx	568 cc.

Fiat lotio. Solve et misce.

Sig.—Utenda si opus sit. P.P.A.

339

R Unguenti Zinci Benzoati	-	ʒij	30 gm.
Spiritus Rectificati	-	ʒij	7 cc

Fiat ung. Misce.

Sig.—Applicandum nocte maneque.

In chronic cases.

(Erasmus Wilson).

340

R Hydrargyri Ammoniati	-	grs. x	.05 gm.
Liquoris Picis Carbonis	-	ʒi	3.5 cc.
Liquoris Plumbi Subacetatis	-	ʒss	1.8 cc.
Vaselinii	-	ʒii	8 gm.
Lanolini ad	-	ʒi	30 gm.

Fiat unguentum. Misce.

Sig.—Applicandum bis in die.

In chronic cases.

341

R Cretæ Preparatæ	-	-	-
Adipis Benzoati	-	ʒā ʒij	30 gm.

Fiat ung. Misce.

(When more stimulating applications irritate).

In chronic cases.

Fœtid Feet—

342

R Acidi Salicylici	-	-	gr. xxx	2 gm.
Pulveris Amyli	-	-	ʒv	18 gm.
Pulveris Talici ad	-	-	ʒiij	90 gm.

Fiat pulvis. Misce.

Sig.—Dust into the socks every morning.

Herpes Zoster—

343

R Potassii Chloratis	-	-	gr. xl	2.6 gm.
Acidi Hydrochlorici Diluti				
Spiritus Chloroformi	-	-		
Extracti Cinchonæ Liquidum	aa	ʒj		3.5.
Aquæ ad	-	-	ʒvj	170 cc.

Fiat mist. Solve et misce.

Sig.—Sumat ʒss ter in die.

344

R Bismuthi Subnitratis	-	-	ʒj	4 gm.
Unguenti Simplicis	-	-	ʒj	30 gm.

Fiat ung. Misce.

Sig.—Applicandum bis in die.

345

R Zinci Oxidi	-	-	ʒiij	12 gm.
Bismuthi Subnitratis	-	-	ʒij	8 gm.
Pulveris Amyli	-	-	ʒss	15 gm.

Fiat pulvis. Misce.

Sig.—To be dusted on the eruption, pro re nata.

Lupus—

346

R Antimonii Chloridi	-	-	
Zinci Chloridi	-	-	30 gm.
Farinae Tritici	-	-	q.s.
Aquæ	-	-	q.s.

Fiat pasta sec. art. Misc.

Sig.—Utatur more dicto.

In ulcerated type.

347

R Potassæ Causticæ	-	-	
Calcis	-	-	30 gm.
Spiritus Rectificati	-	-	q.s.

Fiat pasta. Misc.

Sig.—Utatur more dicto.

In ulcerated type.

348

R Acidi Arseniosi	-	-	gr. xij .8 gm.
Hydrargyri Subchloridi	-	-	30 gm.

Fiat pulvis. Misc.

Sig.—Utatur more dicto.

In ulcerated type.

Requires care. (Dupnytren's Powder).

349

R Acidi Arseniosi	-	-	gr. x .65 gm.
Hydrargyri Persulphatis	-	-	gr. xl 2.6 gm.
Carbonis Animalis	-	-	gr. x .65 gm.

Fiat pulv. Misc.

Sig.—Utatur more dicto.

In ulcerated type.

Pediculosis

350

R Benzolis	-	-	30 cc.
------------	---	---	--------

Sig. Applicandum partibus affectis quotidie.

351

R Acidi Sulphurosi	-	-	℥jv	115 cc.
Aquæ Destillatæ ad	-	-	℥viij	230 cc.
Fiat lotio. Misce.				
Sig.—Applicanda bis in die.				

Psoriasis—

352

R Picis Liquidæ	-	-	℥ij	12 cc.
Fiat pil sec. art. Mitte xii tales. Misce.				
Sig.—Sumat unum ter in die.				

353

R Acidi Hydrocyanici Diluti	-	-	℥xl	2.4 cc.
Olei Cadin	-	-	℥j	30 cc.
Saponis Mollis	-	-	℥ij	60 gm.
Olei Rosmarini	-	-	℥jss	5.3 cc.
Aquæ ad	-	-	℥viij	230 cc.
Fiat lotio. Misce.				

Sig.—To be rubbed over the parts affected night and morning, if the skin is whole.

354

R Acidi Carbolici (Cryst)	-	gr. lx	4 gm.
Unguenti Zinci	-	℥j	30 gm.
Fiat ung. Misce.			
Sig.—Applicandum nocte maneque.			

Pruritis (*see also* 422 to 425)—

355

R Sodii Salicylatis	-	-	gr. ss	1.3 gm
Aquæ ad	-	-	℥ii	60 cc.
Fiat lotio. Misce.				
Sig.—Applicanda nocte maneque.				

356

R Cocainæ Hydrochloridi	-	gr. x	.65 gm.
Adipis Benzoati	-	℥ij	60 gm.
Camphoræ	-	℥j	4 gm.

Fiat unguentum. Misce.

Sig.—Utendum omni nocte.

357

R Atropinæ Sulphatis	-	gr. j	.065 gm.
Boracis	-	gr. cxx	8 gm.
Glycerini	-	℥ss	15 cc.
Acidi Hydrocyanici Diluti	-	℥j	3.5 cc.
Aquæ Aurantii Floris	-	℥ij	60 cc.
Aquæ ad	-	℥xij	340 cc.

Fiat lotio. Solve et misce.

Sig.—Utenda si opus sit.

358

R Acidi Hydrocyanici Diluti	-	℥ij	7 cc.
Plumbi Acetatis	-	gr. lx	4 gm.
Spiritus Rectificati	-	℥j	30 cc.
Aquæ Sambuci ad	-	℥viiij	230 cc.

Fiat lotio. Solve et misce.

Sig.—To be used twice a day. (The skin must not be wounded.)

Purpura (*see also* 87)

359

R Quin. Sulphatis	-	gr. ij	.13 gm.
Tinct. Ferri Perchlor.	-	℥x	.6 cc
Liquoris Arsenici Hydroch-			
lorici	-	℥iiij	.18 cc.
Syrupi Zingiberis	-	℥j	3.5 cc.
Aquæ ad	-	℥jss	45 cc.

Fiat haustus. Mitte vi tales. Solve et misce.

Sig.—Take two or three times a day after food.

Ringworm—

360

R Iodi	5i	4 gm.
Olei Picis Rectificati	5iv	15 cc.
Fiat applicatio. Misce bene.						
Sig.—Applicanda more dicto.						
(Coster's Paste.)						

361

R Acidi Pyrogallici	.	.	.	gr. x	.65 gm.
Sulphuris Precipitati	.	.	.	5ss	2 gm.
Hydrargyri Ammoniaci	.	.	.	gr. xv	1 gm.
Vasellini	.	.	.	5i	30 gm.
Fiat ung. Misce.					
Sig.—Utendum nocte manequ.					

362

R Thymolis	.	.	.	5ss	2 gm.
Chloroformi	.	.	.	5ij	7 cc.
Olei Olive	.	.	.	5vss	20 cc.
Fiat applicatio. Misce.					
Sig.—Utenda quotidie.					

363

R Chrysarobini
Acidi Salicylici	.	.	.	5ā 5i	4 gm.
Paraffini Mollis ad	.	.	.	5ii	60 gm.
Fiat unguentum. Misce.					
Sig.—Applicandum bis in die.					

Scabies—

364

R Sulphuris Sublimati	.	.	.	5j	4 gm.
Balsami Peruviani	.	.	.	5ss	15 gm.
Adipis	.	.	.	5j	30 gm.
Fiat unguentum. Misce.					
Sig.—Utendum mane nocteque.					

365

R Sulphuris Sublimati	-	•	5i	4 gm.
Cretæ Preparatæ	-	•	5ss	2 gm.
Adipis Benzoati ad	-	•	5i	30 gm.

Fiat unguentum. Misce.

Sig.—Applicandum mane nocteque.

366

R Unguenti Sulphuris	-	•	5j	30 gm.
Tinct. Lavandulæ Compositæ	5j			3.5 cc.

Fiat ung. Misce.

Sig.—Utendum omni nocte.

367

R Creosoti	-	•	℥vj	.35 cc.
Unguenti Hydrargyri	-	gr. xxx		2 gm.
Hydrargyri Oxidi Rubri	-	gr. xx		1.3 gm.
Adipis Recentis ad	-	•	5j	30 gm.

Fiat ung. Misce.

Sig.—Applicandum nocte.

368

R Sodii Hyposulphitis	-	•	5ij	8 gm.
Glycerini	-	•	5ss	15 cc.
Aquæ Rosæ ad	-	•	5jv	115 cc.

Fiat lotio. Solve et misce.

Sig.—Applicanda bis terve in die.

Syphilitic Eruptions —

369

R Hydrargyri Iodidi Viridis	-	gr. vij	.8 gm.
Extracti Lupuli	-	gr. lx	4 gm.
Extracti Opii	-	gr. v	.32 gm.

Fiat massa. Divide into twenty-four pills. Misce.

Sig.—Sumat unam ter in die.

370

R Hydrargyri Biniodidi -	-	gr. iij	.2 gm.
Potassii Iodidi -	-	ʒij	8 gm.
Spiritus Rectificati -	-	ʒj	3.5 cc.
Syrupi Zingiberis -	-	ʒjv	15 cc.
Aquæ Distillatæ -	-	ʒjss	45 cc.

Fiat mistura. Solve et misce.

Sig.—Take thirty drops three times a day in a wine-glass of water after food.

371

R Linimenti Opii -	-	ʒvi	21 cc.
Linimenti Belladonnæ -	-		
Linimenti Hydrargyri -	ʒā ʒj		30 cc.

Fiat pigm. Misce.

Sig.—To be used if the eruptions are painful.

372

R Argenti Nitratis -	-	ʒj	4 gm.
Aquæ Distillatæ -	-	ʒj	30 cc.

Fiat pigm. Solve.

Sig.—To be used for discharging sores.

Warts—

373

R Hydrargyri Bichloridi -	-	gr. v	.32 gm.
Acidi Salicylici -	-	ʒj	4 gm.
Collodii ad -	-	ʒj	30 cc.

Fiat pigm. Solve et misce.

Sig.—To be applied every day; the upper crust to be removed daily before each application.

374

R Acidi Salicylici -	-	ʒi	4 gm.
Zinci Chloridi -	-	ʒss	2 gm.
Collodii Flexilis -	-	ʒi	30 cc.

Fiat pigm. Solve et misce.

Sig.—Applicandum nocte.

Wounds (superficial)—

375

R Tinctura Benzoini Composita $\bar{\text{ss}}$ 30 cc.
 Sig.—To be applied on lint and left undisturbed
 for some days.

Ulcers—

376

R Scarlet Red gr. xv 1 gm.
 Unguenti Zinci ad $\bar{\text{ssvii}}$ 25 gm.
 Fiat ung. Misce.

Sig.—Apply on gauze covered with a bandage.
 Use bland ointment to protect surrounding skin.
To promote epithelium formation.

377

R Plumbi Nitratis $\bar{\text{ssj}}$ 4 gm.
 Aquæ ad $\bar{\text{ssxx}}$ 568 cc.
 Fiat lotio. Solve.

Sig.—Apply on lint covered with gutta-percha
 tissue twice a day.
In inflamed ulcers.

378

R Cupri Sulphatis $\bar{\text{ssj}}$ 4 gm.
 Unguenti Simplicis $\bar{\text{ssj}}$ 30 gm.
 Fiat unguentum. Dilute if too strong. Misce.

Sig.—Applicandum omni nocte.
In callous ulcers.

Urticaria—

379

R Liquoris Calcis
 Aquæ Laurocerasi
 Glycerini $\bar{\text{ssss}}$ 30 cc.

Fiat lotio. Misce.
 Sig.—Utenda sæpe.

380

R Hydrargyri Perchloridi	-	gr. jss	.1 gm.
Chloroformi	-	℥℥ _{xx}	1.2 cc.
Glycerini	-	℥ij	60 cc.
Aquæ Rosæ ad	-	℥vüj	230 cc.

Fiat lotio. Solve et misce.

Sig.—To be dabbed on the affected parts.

URINARY DISEASES.

Eladder (irritable)—

381

R Potassii Bicarbonatis	-	℥ji	8 gm.
Tinct. Hyoscyami	-	℥vj	21 cc.
Infusi Buchu ad	-	℥vj	170 cc.

Fiat mistura. Misce.

Sig. ℥j ter in die ex aquæ cyatho sumenda.

382

R Extracti Opii	-	gr. ij	.13 gm.
Extracti Belladonnæ	-	gr. ss	.032 gm.
Olei Theobromatis	-	gr. xij	.8 gm.

Fiat suppositorium. Mitte xii talia. Misce.

Sig.—One to be introduced night and morning.

383

R Zinci Oxidi	-	gr. lx	4 gm.
Extracti Belladonnæ	-	gr. xl	2.6 gm.
Olei Theobromatis	-	℥j	30 gm.
Olei Olivæ	-	℥ij	7 cc.

Fiat massa. Divide into eight pessaries. Misce.

Sig. One to be used when required.

Calculus (urinary)—

384

R Potassii Citratis	•	•	•	•	•	5iij	12 gm.
Aquæ ad	•	•	•	•	•	5iij	90 cc.

Fiat mist. Solve.

Sig.—Take 5ij every three hours in water.

385

R Tincturæ Belladonnæ	•	•	•	•	•	5iss	5.3 cc.
Tincturæ Hyoscyami	•	•	•	•	•	5i	30 cc.
Tincturæ Aurantii	•	•	•	•	•	5iv	15 cc.
Aquæ ad	•	•	•	•	•	5viii	230 cc.

Fiat mistura. Misce.

Sig.—Capiat 5iv quartis horis.

Cystitis—

386

R Spiritus Etheris Nitrosi	•	•	•	•	•	5ij	7 cc.
Spiritus Ammoniæ Aromatici	•	•	•	•	•	5iij	10.5 cc.
Tinct. Lavandulæ Compositæ	•	•	•	•	•	5iss	5.3 cc.
Infusi Buchu ad	•	•	•	•	•	5vi	170 cc.

Fiat mistura. Misce.

Sig.—Capiat cochleare magnum ter in die ex aquâ.

387

R Urotropini	•	•	•	•	•	5ii	8 gm.
Aquæ Menthæ Piperitæ ad	•	•	•	•	•	5ij	60 cc.

Fiat mistura. Solve.

Sig.—Sumat coch. min. vel med. ex aquæ
cyatho quartis horis.*With septic urine.*

388

R Ammonii Benzoatis	•	•	•	•	•	5iv	15 gm.
Tincturæ Lavandulæ Com-	•	•	•	•	•	•	•
positæ	•	•	•	•	•	5vi	21 cc.
Aquæ ad	•	•	•	•	•	5vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat 5iv ter in die.

With alkaline urine.

389

R Acidi Borici Pulveris	℥iii	12 gm.
Glycerini	℥i	30 cc.
Tincturæ Hyoscyami	℥vi	21 cc.
Aquæ ad	℥viii	230 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ℥iv ter in die post cibos. P.P.A.
With acid urine.

390

R Iodoformi (pulv.)	℥ss	15 gm.
Mucilaginis Tragacanthæ	℥ij	60 cc.
Aquæ Destillatæ ad	℥viii	230 cc.

Fiat injectio. Misce.

Sig.—℥ij to ℥iv in half a pint of warm water
as an injection. P.P.A.

Dropsy (renal)—

391

R Elaterii	gr. ¼	.016 gm.
Capsici	gr. ½	.032 gm.
Extracti Colocynthis Co.	gr. 4	.26 gm.

Fiat pilula. Mitte iv tales. Misce.

Sig.—Capiat unam omni mane ante jentaculum.

392

R Potassii Tartratis Acidi	grs. xx	1.3 gm.
Spiritus Juniperi	℥xxx	1.8 cc.
Tincturæ Scille	℥xv	.9 cc.
Tincturæ Digitalis	℥v	.3 cc.
Aquæ Chloroformi ad	℥i	30 cc.

Fiat haustus. Mitte viii tales. Solve et misce

Sig.—Sumat ter in die.

393

R Pulveris Jalapæ Compositi	℥i	4 gm.
---------------------------------------	----	-------

Fiat pulvis. Mitte quattuor tales.

Sig.—Capiendus unus omni mane ante jentaculum
ex aquâ.

Hæmaturia—

394

R Tincturæ Ferri Perchloridi	ʒii	7 cc.
Acidi Phosphorici Diluti.		
Tincturæ Digitalis	- āā ʒiss	5.3 cc.
Glycerini	- - - ʒiv	15 cc.
Aquæ ad	- - - ʒvi	170 cc.
Fiat mistura. Misce.		
Sig.—Capiat ʒiv ter in die.		

Incontinence of Urine—

395

R Potassii Citratis	- - -	
Potassii Bicarbonatis	- āā ʒij	8 gm.
Tincturæ Belladonnæ	- ʒi	3.5 cc.
Syrupi Aurantii	- - - ʒvi	21 cc.
Aquæ ad	- - - ʒvi	170 cc.
Fiat mistura. Solve et misce.		
Sig.—Habeat coch. mag. ex aquā quartis vel sextis horis.		

396

R Spiritūs Etheris Nitrosi	- ʒij	7 cc.
Potassii Citratis	- - - ʒi	4 gm.
Tincturæ Belladonnæ	- ʒi	3.5 cc.
Syrupi Zingiberis	- - - ʒvi	21 cc.
Infusi Buchu ad	- - - ʒvi	170 cc.
Fiat mistura. Solve et misce.		
Sig.—Cap. coch. mag. quartis horis ex cyatho aquæ.		

397

R Phenazoni	- - - ʒi	4 gm.
Syrupi Simplicis	- - - ʒss	15 cc.
Aquæ ad	- - - ʒiiij	90 cc.
Fiat mistura. Solve et misce.		
Sig.—Sum. coch. mag. horā sextā (p.m.) et postea horā nonā.		

Nephritis—

398

R Potassii Citratis	-	-	-	℥iii	12 gm.
Liquoris Ammonii Acetatis				℥iiss	45 cc.
Tincturæ Digitalis	-	-	-	℥iiss	5.3 cc.
Aquæ Chloroformi ad	-	-	-	℥vi	170 cc.

Fiat mistura. Solve et misce.

Sig.—Capiat ℥iv quartis horis.

In the early stages.

399

R Potassii Nitratis	-	-	-	gr. x	.65 gm.
Sodæ Tartarata	-	-	-	gr. xl	2.6 gm.
Aquæ Menthæ Piperitæ ad				℥i	30 cc.

Fiat haustus. Mitte vi tales. Solve.

Sig.—Capiendus bis in die.

In the early stages.

400

R Tinct. Ferri Perchloridi	-			℥ij	7 cc.
Spiritus Ætheris Nitrosi	-			℥jv	15 cc.
Infusi Quassie ad	-	-	-	℥vj	170 cc.

Fiat mistura. Misce.

Sig.—Sumat ℥ss ter in die post cibos.

In second stage.

(Sir G. Stewart.)

401

R Olei Eucalypti	-	-	-	℥i	3.5 cc.
Olei Olivæ	-	-	-		
Olei Ricini	-	-	-	℥ā ℥j	30 cc.

Misce.

For lubricating urethral instruments, catheters, bougies, etc., before use.

UTERINE AND GENITAL DISEASES.

After Pains—

402

R Tinct. Opii -	℥jss	5.3 cc.
Spiritus Ætheris Nitrosi . .	℥ij	7 cc.
Aquæ Camphoræ ad	℥vj	170 cc.
Fiat mistura. Misc.		
Sig. Sumat ℥j quartâquâque horâ.		

403

R Liq. Morphine Hydroch-		
loridi	℥xv	.9 cc.
Spiritus Ammonia Aromatici	℥ss	1.8 cc.
Aquæ Camphoræ ad	℥jss	45 cc.
Fiat haustus. Misc.		
Sig. - Statim sumendus.		

Amenorrhœa—

404

R Ferri et Ammonii Citratis		
Ammonii Bromidi	ââ ℥ij	8 gm.
Decocti Aloes (conc. 1 to 3)	℥ss	15 cc.
Aquæ Mentha Piperita . .	℥viii	230 cc.
Fiat mistura. Solve et misce.		
Sig. Capiat coch. mag. ter in die ex aquâ.		

405

R Tinct. Sanguinaria		
Tinct. Nucis Vomica	ââ ℥ij	7 cc.
Tinct. Aloes	℥ss	15 cc.
Fiant guttae. Misc.		
Sig. - Capiat ℥xx bis in die ex aquâ.		

406

R Hydrargyri Perchloridi . . . gr. $\frac{3}{4}$.05 gm.
 Sodii Arsenatis gr. j .065 gm.
 Ferri Sulphatis Exsiccati . . gr. xxx 2 gm.
 Potassii Carbonatis gr. xv 1 gm.
 Extracti Nucis Vomicae . . gr. v .32 gm.
 Fiat massa. Divide in pilulas triginta. Misce.
 Sig.—One pill before each meal.

407

R Zinci Valerianatis gr. xvij 1.2 gm.
 Pilulae Galbani Compositae gr. xxxvj 2.4 gm.
 Olei Sabinæ ℥ xxxvj 1.6 cc.
 Fiat massa. Divide in pilulas duodecim. Misce.
 Sig.—Take one three times a day.
In hysterical cases.

Cancer (uterine)—

408

R Acidi Salicylici gr. vj .4 gm.
 Sodii Salicylatis ʒij 12 gm.
 Tinct. Eucalypti Gummi . . ʒvj 21 cc.
 Aquæ ad ʒvj 170 cc.
 Fiat injectio. Solve et misce.
 Sig.—Three table-spoonfuls to a pint of water,
 and use as an injection every three or four hours.

409

R Liquoris Zinci Chloridi . . ʒij 7 cc.
 Aquæ ad ʒvij 230 cc.
 Fiat lotio. Solve.
 Sig.—Utenda more dicto.

Cervicitis—

410

R Acidi Benzoici	-	-	-	gr. x	.65 gm.
Collodii	-	-	-	ʒjv	15 cc.
Acidi Tannici	-	-	-	gr. xv	1 gm.
Acidi Carbolici (cryst)	-	-	-	gr. xxv	1.6 cc.
Fiat pig. Solve et misce.					

Sig.—To be applied to the affected part with a small brush through a speculum.

Dysmenorrhœa—

411

R Tinct. Cannabis Indicæ	-	℥℥xv	.9 cc.
Spiritus Juniperi	-	℥℥xxx	1.8 cc.
Spiritus Ætheris	-	℥℥xv	1.2 cc.
Tinct. Aconiti	-	℥℥x	.6 cc.
Misturæ Amygdalæ	-	ʒi	30 cc.
Fiat haustus. Mitte vi tales. Misce.			
Sig.—Capiendus hora somni. P.P.A.			

412

R Tinct. Chlorotormi Com-			
positæ	-	-	ʒij
Potassii Bromidi	-	-	gr. xl
Infusi Taraxaci	-	-	ʒi
Infusi Aurantii ad	-	-	ʒviii
Fiat mistura. Solve et misce.			

Sig.—Capiat coch. mag. duo vel tria ter in die.

413

R Extracti Opii	-	-	-
Extracti Belladonnæ	-	āā gr. jss	.1 gm.
Quinina Sulphatis	-	-	gr. xv
Fiat massa. Divide in pilulas duodecim. Misce.			
Sig.—One to be taken every three hours.			

Leucorrhœa—

414

R Acidi Carbolici Liquifacti	-	3j	3.5 cc.
Glycerini	-	5j	30 cc.
Aquæ Calidæ ad	-	5viij	230 cc.
Fiat injectio. Misce.			

Sig.—To be used with a uterine douche.

415

R Liquoris Plumbi Subacetatis			
Tinct. Opii	-	āā 5j	30 cc.

Fiat lotio. Misce.

Sig.—A tea-spoonful in half-a-pint of warm water, to be injected into the vagina twice daily as a douche.

Menorrhagia—

416

R Liq. Strych. Hydrochloridi	℥iij	.18 cc.
Tinct. Digitalis	-	℥x .6 cc.
Extracti Hydrastis Liq.	-	℥xv .9 cc.
Aquæ ad	-	5j 30 cc.

Fiat haustus. Mitte viii tales. Misce.

Sig. Sumat ter in die.

Nipples (sore)—

417

R Ichthyolis	-	3j	4 gm.
Lanolini	-	3jss	6 gm.
Glycerini	-	3ij	7 cc.
Olei Olivæ	-	3ijss	8.8 cc.

Fiat applicatio. Misce.

Sig.—Apply to nipples, wash off before nursing.

Orchitis—

418

R Morphinae Sulphatis	-	-	gr. j	.065 gm.
Chloralis Hydratis	-	-	ʒjss	6 gm.
Potassii Bromidi	-	-	ʒij	8 gm.
Syrupi	-	-	ʒiv	15 cc.
Aquae ad	-	-	ʒiiij	90 cc.

Fiat mistura. Solve et misce.

Sig.—Sumat ʒss tertiâquâque horâ si opus sit.

419

R Guaiacolis	-	-	-	ʒss	1.8 cc.
Vaselini ad	-	-	-	ʒj	30 cc.

Fiat ung. Misce.

Sig.—Apply to the scrotum on lint twice or thrice daily.

420

R Spiritus Rectificati	-	-	-	-	-
Acidi Acetici Diluti	-	-	ʒi ʒi	30 cc.	
Ammonii Chloridi	-	-	ʒiiij	12 gm.	
Aquae ad	-	-	-	ʒxxx	568 cc.

Fiat lotio. Solve et misce.

Sig.—Apply and allow to evaporate on the testicles.

Ovaritis or Pelvic Cellulitis —

421

R Plumbi Iodidi	-	-	gr. lxxx	5.3 gm.
Extracti Belladonnae	-	-	gr. xxx	2 gm.
Olei Theobromatis	-	-	ʒj	30 gm.

Melt and divide into eight pessaries. Misce.

Sig.—One to be placed every night in the vagina.

Pruritis Vulvæ (*see also* 355 to 358)—

422

R Liquoris Ammonii Acetatis	℥j	30 cc.
Acidi Hydrocyanici Diluti	℥jss	5.3 cc.
Lotionis Plumbi c̄ opio ad	℥viij	230 cc.

Fiat lotio. Misce.

Sig. — Sponge the part twice a day.

423

R Boracis	℥ss	15 gm.
Morphinæ Sulphatis	gr. vj	.4 gm.
Aquæ Rosæ ad	℥viii	230 cc.

Fiat lotio. Solve et misce.

Sig. — Utenda si opus sit.

424

R Acidi Carbolici (cryst)	℥j	4 gm.
Morphinæ Sulphatis	gr. x	.65 gm.
Acidi Borici	℥ij	8 gm.
Vasellini	℥ijss	75 gm.

Fiat unguentum. Misce.

Sig. — Applicandum more dicto.

425

R Camphoræ	gr. lx	4 gm.
Spiritus Rectificati	q.s.	q.s.
Hydrargyri Subchloridi	gr. xl	2.6 gm.
Adipis	℥j	30 gm.

Fiat ung. Misce.

Sig. — Utendum mane nocteque.

Rigidity of Os Uteri—

426

R Antimonii Tartarati	gr. ij	.13 gm.
Tinct. Opii	℞xx	1.2 cc.
Aquæ ad	℥vj	170 cc.

Fiat mist. Solve et misce.

Sig. — Sumat ℥ss omni horâ donec nausea vel vomitus supervenerit.

Subinvolution—

427

R Extracti Ergotæ Liquidi - ʒss 1.8 cc.
 Potassii Chloratis - - gr. x .65 gm.
 Acidi Hydrochlorici Diluti ℥x .6 cc.
 Infusi Gentianæ Co. ad - ʒj 30 cc.
 Fiat haustus. Mitte viii tales. Solve et misce.
 Sig.—Three times a day for three or four months.

Vaginismus—

428

R Iodoformi - - - gr. xv 1 gm.
 Extracti Belladonnæ - - gr. viij .52 gm.
 Olei Theobromatis - - q.s. q.s.
 Divide in sex suppositoria. Misce.
 Sig.—Utendum unum horā decubitus.

PRESCRIPTIONS FOR CHILDREN.

Apthæ or Thrush—

429

R Acidi Borici - - - ʒi 4 gm.
 Thymolis - - - gr. i .065 gm.
 Glycerini - - - ʒi 30 cc.
 Fiat pigmentum. Solve.
 Sig.—Applicandum ope penicilli vel spongiolæ
 pro re nata.

Bronchitis—

430

R Vini Ipecacuanhæ	. . .	℥xv	.9 cc.
Spiritus Ammoniac Aromatici		℥v	.3 cc.
Syrupi Tolutani	. . .	ʒiij	10.5 cc.
Aquæ Anethi ad	. . .	ʒjss	45 cc.

Fiat mistura. Misce.

Sig.—Detur ʒj ter in die.

(At 6 months old.)

431

R Vini Ipecacuanhæ	. . .		
Spiritus Ætheris Nitrosi		āā ʒj	3.5 cc.
Liquoris Ammonii Acetatis		ʒiss	5.3 cc.
Syrupi Aurantii	. . .	ʒiij	10.5 cc.
Aquæ Anethi ad	. . .	ʒjss	45 cc.

Fiat mist. Misce.

Sig.—Habeatur ʒj quartâquâque horâ.

(At 1 year old.)

432

R Vini Ipecacuanhæ	. . .	℥x	.6 cc.
Oxymellis Scillæ	. . .	℥xl	2.4 cc.
Spiritus Ætheris Nitrosi	. . .	℥xx	1.2 cc.
Tinct. Camphoræ Co.	. . .	℥xv	.9 cc.
Aquæ Anisi	. . .	ʒjss	45 cc.

Fiat mistura. Misce.

Sig.—ʒij ter in die habendæ.

(At 1 year old.)

433

R Infusi Senegæ	. . .	ʒj	30 cc.
Liquoris Ammonii Acetatis		ʒss	15 cc.
Syrupi Scillæ	. . .		
Syrupi Papaveris	. . .	āā ʒij	7 cc.

Fiat mistura. Misce.

Sig.—Detur ʒj ter in die.

(At 2 years.)

434

R Ammonii Carbonatis	.	.	gr. xii	.8 gm.
Tinct. Scillæ	.	.	℥xvj	1 cc.
Syrupi Tolutani	.	.	ʒiij	10.5 cc.
Infusi Senegæ ad	.	.	ʒiij	90 cc.

Fiat mistura. Solve et misce.

Sig.—Dentur ʒij quartâquâque horâ.

(At 2 or 3 years old.)

Catarrh—

435

R Tinct. Camphoræ Co.	.	.	℥xx	1.2 cc.
Vini Ipecacuanhæ	.	.	℥x	.6 cc.
Vini Antimonialis	.	.	℥xxx	1.8 cc.
Misturæ Amygdalæ ad	.	.	ʒj	30 cc.

Fiat mistura. Misce.

Sig.—Dentur ʒij quartâquâque horâ. P. P. A.

(At 1 year old.)

436

R Vini Ipecacuanhæ	.	.	ʒij	7 cc.
Mucilaginis Acaciæ	.	.	ʒss	15 cc.
Syrupi Limonis	.	.	ʒj	3.5 cc.
Syrupi Papaveris	.	.	ʒjss	5.3 cc.
Aquæ ad	.	.	ʒii	60 cc.

Fiat mist. Misce.

Sig.—One to two teaspoonfuls to be given every three hours.

(At 3 years.)

437

R Vini Antimonialis	.	.	mxlv	2.7 cc.
Vini Ipecacuanhæ	.	.	ʒj	3.5 cc.
Syrupi Rhæados	.	.	ʒiij	10.5 cc.
Liquoris Ammonii Acetatis	.	.	ʒiv	15 cc.
Aquæ ad	.	.	ʒiij	60 cc.

Fiat mist. Misce.

Sig.—Dentur drachmæ duæ quartis horis.

(At 2 or 3 years old.)

Chorea—

438

R Tincturæ Ferri Perchloridi
 Liq. Arsenici Hydroch-
 lorici ʒi ℥xv .9 cc.
 Glycerini ʒiv 15 cc.
 Aquæ Menthæ Piperitæ ad ʒij 60 cc.
 Fiat mistura Misce.
 Sig.—Detur coch. minimum ter in die ex aquâ
 post cibum.

439

R Liquoris Arsenicalis ℥xv .9 cc.
 Ammonii Bromidi ʒi 4 gm.
 Syrupi Aurantii ʒi 30 cc.
 Aquæ Chloroformi ad ʒii 60 cc.
 Fiat mistura. Solve et misce.
 Sig.—Detur coch. parvum bis in die post
 cibum ex aquâ.

Convulsions—

440

R Sodii Bromidi gr. iv .26 gm.
 Phenazoni gr. ij .13 gm.
 Syrupi Simplicis ʒi 30 cc.
 Aquæ Chloroformi ad ʒii 60 cc.
 Fiat mist. Solve et misce.
 Sig.—Detur drachma tertiis horis si opus sit.

Constipation—

441

R Pulveris Rhei gr xv 1 gm.
 Magnesii Carbonatis Levis gr. lx 4 gm.
 Aquæ Anethi ad ʒjss 45 cc.
 Fiat mist. Misce.
 Sig.—A teaspoonful to be given every two
 or three hours in milk.
 (At 12 months old.)

442

- R Hydrargyri cum Cretâ - - -
 Pulveris Rhei - - - āā gr. j .065 gm.
 Sodii Bicarbonatis - - - gr. ij .13 gm.
 Fiat pulvis. Mitte xii tales. Misce.
 Sig.—Unus nocte si opus sit habendus.
 (At 1 year old.)

443

- R Magnesii Sulphatis - - - ʒi 4 gm.
 Tinct. Rhei - - - ʒij 7 cc.
 Syrupi Zingiberis - - - ʒj 3.5 cc.
 Aquæ Carui ad - - - ʒjss 45 cc.
 Fiat mist. Solve et misce.
 Sig.—Detur ʒj ter in die.

444

- R Pulveris Rhei - - -
 Pulveris Scammonii - - -
 Potassii Sulphatis - - - āā gr. x .65 gm.
 Pulveris Cinnamomi Com-
 positi - - - gr. vj .4 gm.
 Tere optime simul. Divide in pulveres sex.
 Sig.—Sumat unum quartâquâque horâ donec
 alvus soluta sit.

445

- R Magnesii Sulphatis - - - ʒij 8 gm.
 Syrupi Aurantii - - - ʒij 7 cc.
 Aquæ Carui ad - - - ʒjss 45 cc.
 Fiat mist. Solve et misce.
 Sig.—Give ʒij every hour till the bowels act.
 (At 3 years old.)

Croup—

446

- R Vini Ipecacuanhæ - - - ʒj 3.5 cc.
 Sig.—To be given at once, and followed by a
 teacupful of warm water as an emetic.

447

R Oxymellis Scillæ	-	-	ḡij	30 cc.
Vini Ipecacuanhæ	-	-	ḡij	60 cc.

Fiat mist. Misce.

Sig.—Give a teaspoonful every 10 or 15 minutes till sickness is produced.

448

R Potassii Bicarbonatis	-	-	gr. xl	2.6 gm.
Acidi Citrici	-	-	gr. xx	1.3 gm.
Vini Antimonialis	-	-	ḡjss	5.3 cc.
Vini Ipecacuanhæ	-	-	℥xx	1.2 cc.
Syrupi Limonis	-	-	ḡjss	8.8 cc.
Aquæ ad	-	-	ḡiij	90 cc.

Fiat mist. Solve et misce

Sig.—Give ḡij every three or four hours.

(At 2 years old, after the child has been placed in a hot bath.)

Debility—

449

R Ferri et Ammonii Citratis	-	ḡi	4 gm.
Spiritus Ammoniac Aromatici	-	-	-
	-	ḡjss	5.3 cc.
Tincture Cinchonæ Co.	-	ḡi	3.5 cc.
Syrupi Limonis	-	-	-
	-	ḡss	15 cc.
Aquæ ad	-	-	-
	-	ḡiij	90 cc.

Fiat mist. Misce.

Sig.—Detur drachma ter in die post cibos.

Dentition

450

R Cocainæ Hydrochloridi	-	gr. jss	.1 gm.
Tinct. Conii	-	-	.
Syrupi	-	āā ʒij	7 cc.

Fiat lotio. Solve et misce.

Sig.—Rub on the gums several times daily.

Diarrhœa—

451

R Misturæ Cretæ	-	-	-	5vj	22 gm.
Pulveris Cretæ Aromatici	-	-	-	5j	4 gm.
Extracti Opii Liq.	-	-	-	℥xv	.9 cc.
Aquæ Cinnamomi ad	-	-	-	5jv	115 cc.

Fiat mist. Misce.

Sig.—Give a teaspoonful every hour. P.P.A.
(At 1 year old.)

452

Tinct. Catechu	-	-	-	5ij	7 cc.
Syrupi	-	-	-	5iij	10.5 cc.
Decocti Hamatoxyli ad	-	-	-	5jss	45 cc.

Fiat mist. Misce.

Sig.—Detur 5j ter in die.
(At 1 year old.)

453

R Hydrargyri Subchloridi	-	gr. ½	.016 gm.
Bismuthi Carbonatis	-	gr. ii	.13 gm.
Sacchari Lactis	-	gr. iii	.2 gm.

Fiat pulvis. Mitte sex tales. Misce.

Sig.—Unus bis in die habendus.
(At 1 year old.)

454

R Olei Ricini	-	-	-	5j	3.5 cc.
Pulveris Acaciæ	-	-	-	5ss	2 gm.
Sacchari Albi	-	-	-	5ss	2 gm.
Tinct. Opii	-	-	-	℥jv	.24 cc.
Aquæ Cinnamomi ad	-	-	-	5j	30 cc.

Fiat mist. Misce.

Sig.—Detur 5j tertia vel quartaquaque horâ. P.P.A.
(At 1 year old.)

With enteric inflammation.

455

R Acidi Carbolici (cryst)	-	gr. ij	.13 gm.
Bismuthi Subnitratis	-	ʒj	4 gm.
Mucilaginis Acaciæ	-	ʒss	15 cc.
Aquæ Menthæ Piperitæ ad	-	ʒij	60 cc.

Fiat mist. Misce.

Sig.—Give half a tea-spoonful every two or four hours. P.P.A.

(At 1 or 2 years old.)

With offensive motions.

456

R Salolis	-	gr. viii	.52 gm.
Pulv. Tragacanthæ Com-	-		
positæ	-	gr. x	.65 gm.
Bismuthi Salicylatis	-	gr. xx	1.3 gm.
Syrupi Zingiberis	-	ʒiv	15 cc.
Aquæ Chloroformi ad	-	ʒij	60 cc.

Fiat mist. Misce.

Sig.—Detur coch. min. quartis horis. P.P.A.

(At 3 years old.)

With flatulence.

Dyspepsia —

457

R Sodii Bicarbonatis	-	gr. xxjv	1.5 gm.
Tinct. Rhei	-	ʒj	3.5 cc.
Extracti Taraxaci	-	gr. xl	2.6 gm.
Infusi Calumbæ	-	ʒi	30 cc.
Aquæ Carni ad	-	ʒij	60 cc.

Misce.

Sig.—Detur ʒij bis in die.

With acidity.

458

R Liquoris Hydrargyri Perch-		
loridi	℥ss	1.8 cc.
Extracti Taraxaci Liquidi	℥i	3.5 cc.
Extracti Cinchonæ Liquidi	℥ss	1.8 cc.
Tinct. Aurantii	℥j	3.5 cc.
Syrupi	℥iij	10.5 cc.
Aquæ ad	℥iij	60 cc.

Misce.

Sig.—Dentur drachmæ duæ bis die.

(At 3 to 4 years old.)

(West.)

With biliousness.

Favus, Ringworm, etc—

459

R Sodii Hyposulphitis	℥i	30 gm.
Glycerini	℥iii	10.5 cc.
Aquæ ad	℥x	280 cc.

Fiat lotio. Solve et misce.

Sig.—Applicanda mane nocteque affectis partibus.

Fever—

460

R Liquoris Ammonii Acetatis	℥ij	7 cc.
Spiritus Ammoniaë Aromatici	℥vj	35 cc.
Vini Ipecacuanhæ	℥ss	1.8 cc.
Syrupi Aurantii	℥iij	10.5 cc.
Aquæ ad	℥jss	45 cc.

Fiat mist. Misce.

Sig.—Detur ℥j ter in die.

(At 6 to 12 months old.)

461

R Potassii Nitratis	-	-	gr. xij	.8 gm.
Magnesii Sulphatis	-	-	ʒj	4 gm.
Syrupi Limonis	-	-	ʒij	10.5 cc.
Aquæ ad	-	-	ʒjss	45 cc.

Fiat mist. Solve et misce.

Sig.—Dentur ʒij ter in die.

(At 1 year old.)

462

R Vini Ipecacuanhæ	-	-	ʒj	3.5 cc.
Spiritus Ætheris Nitrosi	-	-	ʒjss	5.3 cc.
Potassii Nitratis	-	-	ʒss	2 gm.
Syrupi Aurantii	-	-	ʒss	15 cc.
Aquæ ad	-	-	ʒjss	45 cc.

Fiat mist. Solve et misce.

Sig.—Detur ʒj ter in die.

(At from 1 to 2 years old.)

463

R Potassii Bitartratis	-	-	ʒss	15 gm.
Syrupi Limonis	-	-	ʒii	60 cc.
Aquæ ad	-	-	ʒxii	340 cc.

Fiat mist.

Sig.—Give as a drink.

Flatulence—

464

R Sodii Bicarbonatis	-	-	gr. x	.65 gm.
Spiritus Ammoniac Aromatici	℥	xii		.7 cc.
Syrupi Zingiberis	-	-	ʒij	7 cc.
Aquæ Anethi ad	-	-	ʒjss	45 cc.

Fiat mist. Solve et misce.

Sig.—Detur ʒj pro re nata.

(At a week or more old.)

465

R Sodii Carbonatis	-	-	gr. xij	.8 gm.
Spiritus Ammoniae Aromatici	℥xii			.7 cc.
Syrupi Simplicis	-	-	℥ij	7 cc.
Tinct. Cardamomi Compositae	℥xx			1.2 cc.
Aquae Chloroformi ad	-	-	℥jss	45 cc.

Fiat mist. Solve et misce.

Sig.- Detur ℥j quartâquâque horâ.

(At 6 months old.)

Marasmus—

466

R Syrupi Ferri Phosphatis				
Co.	-	-	℥ij	7 cc.
Aquae ad	-	-	℥jss	45 cc.

Fiat mist. Misce.

Sig.- Detur ℥j ter in die.

(At 1 year old.)

Incontinence of Urine—

467

R Tinct. Belladonnae	-	-		
Tinct. Cubebae	-	-	℥ā ℥ij	7 cc.
Tinct. Nucis Vomicae	-	-		
Tinct. Rhus Aromaticae	℥ā ℥j			3.5 cc.
Tinct. Cascarillae	-	-	℥ij	7 cc.

Fiant. guttae. Misce.

Sig.- Give twelve drops in water at bedtime.

(At from 7 to 10 years.)

Insomnia—

468

R Ammonii Bromidi	-	-	gr. xvi	1 gm.
Tinctura Hyoscyami	-	-	℥ss	1.8 cc.
Syrupi Papaveris Rubri	-	-	℥ii	7 cc.
Glycerini	-	-	℥iii	10.5 cc.
Aquae Chloroformi ad	-	-	℥ii	60 cc.

Fiat mistura. Solve et misce.

Sig.—Detur cochleare minimum hora somni.

(At 2 years old.)

469

R Tinct. Opii	-	-	-	℥j	.06 cc.
Mucilaginis Acaciæ	-	-	-	-	-
Syrupi Simplicis	-	-	-	āā ʒij	7 cc.
Aquæ Anethi ad	-	-	-	ʒj	30 cc.

Fiat mist. Misce.

Sig.—Detur ʒj bis terve in die. P.P.A.

(At 1 year old.)

470

R Pulveris Ipecacuanhæ Com-					
positi	-	-	-	gr. j	.065 gm.
Sacchari Albi	-	-	-	gr. xx	1.3 gm.

Fiat pulv. Divide in iv chartas. Misce.

Sig.—Habeatur unus pro re nata.

(At from 1 to 4 weeks old.)

Œdema of Lung—

471

Pilocarpinæ Nitratis	-	-	gr. ½	.032 gm.
Aquæ Destillatæ	-	-	ʒj	3.5 cc.

Solve.

Sig.—Five minims hypodermically.

(Used with great success in the case of a child 3 years old.)

(A. Weber.)

Pneumonia—

472

R Vini Ipecacuanhæ	•	-	ʒjss	5.3 cc.
Syrupi Aurantii	•	-	ʒss	15 cc.
Tinct. Camphoræ Com-				
positæ	•	•	•	ʒiiij
Liquoris Ammonii Citratis			ʒss	15 cc.
Aquæ ad	•	•	•	ʒij
				60 cc.

Fiat mist. Misce.

Sig.—Sumatur ʒj secunda vel tertiâquâque horâ.

(At from 3 to 5 years old.)

Early stage.

Phthisis—

473

R Acidi Nitrici Diluti	• • •	℥℥xvj	1 cc.
Acidi Hydrochlorici Diluti	• • •	℥℥xxjv	1.4 cc.
Acidi Hydrocyanici Diluti	• • •	℥℥v	.3 cc.
Spiritus Chloroformi	• • •	℥℥xl	2.4 cc.
Syrupi Aurantii	• • •	ʒiv	15 cc.
Aquæ ad	• • •	ʒjv	115 cc.

Fiat mist. Misc.

Sig.—Detur ʒss ter in die.

(At 4 years old.)

With cod-liver oil.

Tabes Mesenterica—

474

R Extracti Taraxaci Liquidi	ʒij	7 cc.
Sodii Bicarbonatis	• • • ʒj	4 gm.
Extracti Sarsæ Liquidi	• • • ʒjv	15 cc.
Syrupi Aurantii	• • • ʒvi	21 cc.
Liquoris Sarsæ Compositi		
Conc. ad	• • • ʒvj	170 cc.

Fiat mist. Solve et misce.

Sig.—Sumatur ʒss ter die in lacte.

(At 4 years old.)

(West.)

475

R Olei Morrhuæ	• • • ʒiijss	105 cc.
Spiritus Ammoniac Aromatici	ʒj	30 cc.
Tinct. Opii	• • • ʒjv	15 cc.
Olei Cajuputi	• • • ʒi	30 cc.

Misc.

Sig.—To be rubbed over the abdomen when cod-liver oil cannot be borne by the stomach.

Whooping Cough—

476

R Ammonii Bromidi	-	-	gr. xij	.8 gm.
Vini Ipecacuanhæ	-	-	ʒj	3.5 cc.
Succi Conii	-	-	ʒss	1.8 cc.
Syrupi Tolutani	-	-	ʒiij	10.5 cc.
Aquæ ad	-	-	ʒjss	45 cc.

Fiat mist. Solve et misce.

Sig.—Sumatur ʒj ter in die.

(At from 1 to 2 years old.)

477

R Phenazoni	-	-	gr. xx	1.3 gm.
Vini Ipecacuanhæ	-	-	ʒi	3.5 cc.
Syrupi Tolutani	-	-	ʒiij	10.5 cc.
Glycerini	-	-	ʒii	7 cc.
Aquæ Chloroformi ad	-	-	ʒii	60 cc.

Fiat mist. Solve et misce.

Sig.—Detur cochleare minimum sextis horis
ex aquâ.

(At 2 to 4 years old.)

478

R Succi Conii	-	-	ʒj	3.5 cc.
Acidi Hydrocyanici Diluti	-	-	℥vi	.35 cc.
Vini Ipecacuanhæ	-	-	ʒj	3.5 cc.
Syrupi Aurantii	-	-	ʒss	15 cc.
Aquæ ad	-	-	ʒjss	45 cc.

Fiat mist. Misce.

Sig.—Sumatur ʒj quartâquâque horâ.

(At 2 years old.)

479

R Potassii Bromidi	-	-	gr. xxiv	1.5 gm.
Tinct. Belladonnæ	-	-	℥xxv	1.5 cc.
Syrupi	-	-	ʒss	15 cc.
Aquæ ad	-	-	ʒjss	45 cc.

Fiat mist. Solve et misce.

Sig.—Detur ʒj ter in die.

(At 2 years old.)

480

R Bromoformi-	-	-	℥x	.6 cc.
Alcoholis	-	-	ʒj	3.5 cc.
Syrupi Simplicis	-	-	ʒiij	10.5 cc.
Aquæ ad	-	-	ʒiij	90 cc.

Fiat mist. Misce.

Sig.—Give one or two tea-spoonfuls daily. P.P.A.

481

R Zinci Sulphatis	-	-	gr. viii	.52 gm.
Tincturæ Belladonnæ	-	-	℥xxx	1.8 cc.
Syrupi Aurantii	-	-	ʒiv	15 cc.
Aquæ ad	-	-	ʒiv	115 cc.

Fiat mistura. Solve et misce.

Sig.—Detur semi-drachma ter in die ex aqua.

(At 3 years old.)

482

R Olei Succini	-	-	-	
Olei Olivæ	-	-	ʒā ʒss	15 cc.
Olei Caryophylli	-	-	ʒj	30 cc.

Fiat linimentum. Misce.

Sig.—To be rubbed down the spine
night and morning.

(A substitute for Roche's embrocation.)

Worms—

483

R Santonini	-	-	ʒii	8 gm.
Saccharini Solubilis	-	-	gr. ii	.13 gm.
Pulveris Scammonii	-	-	ʒii	8 gm.
Sodii Bicarbonatis	-	-	ʒss	2 gm.
Sacchari Lactis ad	-	-	ʒii	60 gm.

Fiat pulvis. Misce.

Sig.—One grain for each year of life up to 5, and
half a grain for each year after. Not to be given
under six months. To be given in syrup or milk.*For round or thread worms.*

484

R Santonini	-	-	gr. ij	.13 gm.
Sacchari Lactis	-	-	gr. x	.65 gm.

Misce.

Sig.—To be given fasting, and followed by a brisk purge.

(At 3 years old.)

For round worms.

485

R Hydrargyri Subchloridi	-	gr. ij	.13 gm.
Pulveris Scammonii Co.	-	gr. jv	.26 gm.
Pulveris Cretæ Aromat.	-	gr. v	.32 gm.

Fiat pulvis. Mitte vi tales. Misce.

Sig.—Unus hora somni sumendus.

(At 4 years old.)

For thread worms.

486

R Pulveris Jalapæ Compositi	-	-	-
Potassii Tartratis Acidi	-	-	-
Ferri Carbonatis	-	āā 5j	30 gm.
Pulveris Zingiberis	-	3ss	2 gm.
Mellis Depurati	-	q.s.	-

Fiat electuarium. Misce.

Sig.—Capiatur 3ss vel 3i nocte maneque.

(At 5 years old.)

For thread worms.

487

R Extracti Filicis Liquidi	-	3ss	1.8 cc.
Pulveris Acaciæ	-	gr. x	.65 gm.
Spiritus Myristicæ	-	℥v	3 cc.
Syrupi Tolutani	-	3jv	15 cc.
Aquæ Cinnamomi ad	-	5j	30 cc.

Fiat haustus. Misce.

Sig. To be given at night followed by a saline purge in the morning.

(At 8 or 10 years old.)

For tape worms.

APPENDIX.

WEIGHTS AND MEASURES.

Imperial Standard.

MEASURES OF MASS.

1 grain	-	-	gr.	
1 ounce (avoir.)	oz.	=	437.5 gr.	
1 pound	lb.	=	16 ounces	= 7000 gr.

The drachm (ʒ) where used as a weight is to be taken as meaning 60 grains, and the ounce (℥) as meaning 480 grains.

MEASURES OF CAPACITY.

1 minim	-	min. or ℥.	
1 fluid drachm	fl. dr. or ʒ	=	60 minims.
1 fluid ounce	fl. oz. or ℥	=	8 fluid drachms.
1 pint	-	O	= 20 fluid ounces.
1 gallon	-	C	= 8 pints.

Metric Weights and Measures and
their Equivalents in the British
Pharmacopœia.

1 gramme	-	gm.	= 15.4323554 grs.
1 centigramme	-	cgm.	= 0.154323 grs.
1 milligramme	-	mgm.	= 0.0154342 grs.
1 litre	-	-	= 35.196 fl. ozs.
1 cubic centimetre	-	cc	= 16.95 minims.
1 metre	-	-	= 39.37 inches

Table for converting Imperial into approximate Metric Standard.

EQUIVALENT MEASURES OF MASS (MARTINDALE).

1 pound	-	-	-	=	453.59 gm.
$\frac{1}{2}$ pound	-	-	-	=	225.8.
$\frac{1}{4}$ pound	-	-	-	=	113.4.
1 ounce	-	-	-	=	30 (28.35).
$\frac{1}{2}$ ounce	-	-	-	=	15.
2 drachms	-	120 gr.	=	8.	
$1\frac{1}{2}$ drachms	-	90 gr.	=	6	
1 drachm	-	60 gr.	=	4.	
2 scruples	-	40 gr.	=	2.6.	
$\frac{1}{2}$ dr.	-	30 gr.	=	2	
24 grains	-	-	-	=	1.5.
1 scruple	-	20 gr.	=	1.3.	
15 grains	-	-	-	=	1
12 grs.	-	-	-	=	.8.
10 grs.	-	-	-	=	.65.
9 grs.	-	-	-	=	.6.
3 grs.	-	-	-	=	.52.
7 grs.	-	-	-	=	.46.
6 grs.	-	-	-	=	.4
3 grs.	-	-	-	=	.2.
2 grs.	-	-	-	=	.13.
1 gr.	-	-	-	=	.065.
$\frac{3}{4}$ gr.	-	-	-	=	.05.
$\frac{1}{2}$ gr.	-	-	-	=	.032.
$\frac{1}{4}$ gr.	-	-	-	=	.02.
$\frac{1}{8}$ gr.	-	-	-	=	.016.
$\frac{1}{16}$ gr.	-	-	-	=	.013.
$\frac{1}{32}$ gr.	-	-	-	=	.01
$\frac{1}{64}$ gr.	-	-	-	=	.008.
$\frac{1}{128}$ gr.	-	-	-	=	.0065.
$\frac{1}{256}$ gr.	-	-	-	=	.0054.
$\frac{1}{512}$ gr.	-	-	-	=	.004.
$\frac{1}{1024}$ gr.	-	-	-	=	.0032.
$\frac{1}{2048}$ gr.	-	-	-	=	.0026.

$\frac{1}{36}$ gr.	-	-	-	=	.0022.
$\frac{1}{32}$ gr.	-	-	-	=	.002.
$\frac{1}{40}$ gr.	-	-	-	=	.0016.
$\frac{1}{50}$ gr.	-	-	-	=	.0013.
$\frac{1}{64}$ gr.	-	-	-	=	.001.
$\frac{1}{100}$ gr.	-	-	-	=	.00065.

EQUIVALENT MEASURES OF CAPACITY.

1	gallon	-	Ci	=	4 55 litres.
1	pint	+ 15 fl. oz.	=	1 litre.	
1	pint	- $\frac{1}{5}$ xx	Oi	=	.57 litres (563 cc).
12	ounces	-	-	=	340 cc.
10	ozs.	-	-	=	280 cc.
8	ozs.	-	-	=	230 cc.
6	ozs.	-	-	=	170 cc.
5	ozs.	-	-	=	140 cc.
4	ozs.	-	-	=	115 cc.
3	ozs.	-	-	=	90 cc.
2	ozs.	-	-	=	60 cc.
1	oz.	-	-	=	30 cc.
$\frac{1}{2}$	oz.	-	(5iv)	=	15 cc.
3	fluid drachms	-	-	=	10.5 cc.
2	fluid drachms	-	-	=	7 cc.
100	minims	-	-	=	6 cc.
90	m.	-	(5iss)	=	5.3 cc.
60	m.	-	(5i)	=	3.5 cc.
50	m.	-	-	=	3 cc.
40	m.	-	-	=	2.4 cc.
30	m.	-	(5ss)	=	1.7 cc.
25	m.	-	-	=	1.5 cc.
20	m.	-	-	=	1.2 cc.
17	m.	-	-	=	1 cc.
15	m.	-	-	=	.9 cc.
12	m.	-	-	=	.7 cc.
10	m.	-	-	=	.6 cc.
8	m.	-	-	=	.5 cc.
6	m.	-	-	=	.35 cc.
5	m.	-	-	=	.3 cc.
4	m.	-	-	=	.24 cc.

3	m.	-	-	-	=	.18 cc.
2	m.	-	-	-	=	.12 cc.
1	m.	-	-	-	=	.06 cc.

Table of Strength of Solution in an Imperial Pint of Water.

1½	grs.	-	-	make	1%	-	-	or 1 in 5,000.
8½	grs.	-	-	"	1½%	-	-	" 1,000.
43½	grs.	-	-	"	½%	-	-	" 200.
87½	grs.	-	-	"	1%	-	-	" 100.
437½	grs.	-	-	"	5%	-	-	" 20.
875	grs.	-	-	"	10%	-	-	" 10.

INCOMPATIBILITIES.

The use of compound drugs is complicated by chemical, pharmaceutical, and therapeutical incompatibilities between many of them. The first two types are more readily ascertained than the third, because they are usually exhibited by alteration in condition, taste, smell, or colour, whilst the latter can only be discovered by the effects of its administration. For the study of the latter, the student is referred to the text-books on *Materia Medica* and *Therapeutics*. The chief chemical and pharmacological incompatibilities might be briefly summarised as follows:—

Mineral acids or acid salts with alkalis or alkaline salts, iodides and bromides, spt. eth. nitrosi, metallic oxides, glucosides, hydrates, tannin

Alkalis and their oxides with acids and acid salts, metallic salts, alkaloids and their salts, chloral, quinine, iodides and bromides, tinct. card. co., and tinct. cocci.

- Iron salts and solutions with tannin-containing drugs, salicylates, carbolic acid, calomel.
- Alkaloids and their salts with alkalies, tannic, phosphoric, and boracic acids, iodides and bromides.
- Mucilage with concentrated alcohol, iron and lead salts, mineral acids nitrous ether, glucosides.
- Nitrous ether with phenazonum, bromides and iodides, iron salts, guaiacum, gallic and tannic acids, mucilages, mineral acids.
- Quinine with alkalies and carbonates, salicylates, astringent infusions.
- Tannic acid with alkaloids, mineral acids, alkalies, antimony, lead, silver and iron salts, arsenic, gelatin, and albumen.
- Nitrate of silver with alkalies and carbonates, chlorides, iodides, arsenic, astringent infusions.
- Perchloride of mercury with vegetable infusions containing tannin or albumen, alkalies, iodides, tartar emetic.
- Calomel with alkalies, iodides, iron and lead salts, prussic acid.
- Iodine and iodides with alkaloids, mineral acids, nitrous ether, metallic salts and starch, e.g., iodide of potassium with preparations of strychnia.
- Lime water with mercurial salts, carbonates, morphine, quinine, and phosphoric acid.
- Carbonates with alkaloids, metallic salts, and acetic acid.
- Nitric acid with alcohol, sulphate of iron, and acetate of lead.
- Phosphoric acid with alkaloids, metallic salts, and lime water.
- Alum with alkalies, tannin, tartrates, and salts of lead.

Cyanides with chloral; cocaine with borax; oxide of silver with creosote; bismuth subnitrate with carbonates and bicarbonates in watery solution; infusion of roses with alkalies: and it should be remembered that chlorate and permanganate of potassium should not be rubbed up with charcoal, sulphur, glycerine, etc., as explosive mixtures may be produced.

Note.—In some cases incompatibility between drugs has been purposely ignored as in the combination of calomel with lime-water (black wash), of hydrochloric acid and chlor. of pot. (chlorine gargle), of iod. of pot. and perchlor. of merc. and in efferv. mixtures. It may also be overcome in many instances by skilful dispensing, *e.g.* by the use of dil. phos. acid in preventing the inky colour of iron mixtures: of liq. ammon. acet. in avoiding the precipitation of quinine from its ammoniated tinct. and of salicylic acid when mixed with water; of liq. ammonize in making liquid extr. of casara soluble: of carb. of pot. in rendering nitrous ether compatible with iod. of pot.; and of the tincts. of senega and quillaria in emulsifying santol, copaiba, and other oils and making them miscible in water.

NUMERALS COMMONLY USED IN PRESCRIPTIONS.

1. Unus-a-um. 1st. Primus-a-um. Once, semel.
2. Duo-re-o. 2nd. Secundus-a-um. Twice, bis.
3. Tres-es-ia. 3rd, Tertius-a-um. Thrice, ter.
4. Quattuor or quatuor. 4th, Quartus-a-um. Four times, quater.
5. Quinque. 5th, Quintus-a-um.

6. Sex. 6th, Sextus-a-um.
 7. Septem. 7th, Septimus-a-um.
 8. Octo. 8th, Octavus-a-um.
 9. Novem. 9th, Nonus-a-um.
 10. Decem. 10th, Decimus-a-um.
 11. Undecim. 11th, Undecimus-a-um.
 12. Duodecim. 12th, Duodecimus-a-um.
 15. Quindecim 16. Sedecim. 18 Duodeviginti.
 20. Viginti. 24. Quattuor et viginti. 25. Quinque et viginti.
 30. Triginta. 40. Quadraginta. 50. Quinquaginta.
 60. Sexaginta. 70. Septuaginta. 80. Octoginta.
 90 Nonaginta.
 100. Centum. 500. Quingenti-æ-a. 1000. Mille (Pl Millia).

EXAMPLES—

Horā unā. One o'clock.

Mane primo. The first thing, or early in the morning.

Semel in die. Once a day.

Cochlearia duo. Two tablespoonfuls.

Secundis horis. Every two hours. Horā secundā. Two o'clock.

Bis in die, bis die, or bis de die. Twice a day.

Grana tria. Three grains.

Tertiis horis. Every three hours. Horā tertiā. Three o'clock.

Ter in die, ter die, or ter de die. Thrice a day.

Minima quattuor. Four minims.

Quartis horis. Every four hours. Horæ pars quarta. A quarter of a hour.

Quater in die. Four times a day.

Unciæ quinque. Five ounces.

Horā quintā. Five o'clock. Quintis horis. Every five hours.

Mitte sex tales. Send six of such.

Sextis horis. Every six hours. Sextā horā. Six o'clock.

TABLE OF QUANTITIES.

Pondera et Mensuræ.

*(Used in the Accusative after an Active Verb,
e.g., Recipe.)*

Cochleare-is. n. 3=a spoonful.

Coch. magnum, amplum, largum, or plenum=
a tablespoonful.

Coch. medium or modicum=a dessertspoonful.

Coch. minimum or parvum=a teaspoonful.

Cochlearia duo, tria, etc.=two, three, etc.,
spoonfuls.

Congius-ii. m. 2=a gallon (Ci).

Cyathus-i m. 2=a glass or cup.

Cyathus amplus or magnus=a tumblerful.

Cyathus parvus or vinosus=a wineglassful.

Dimidius-a-um=half (adj). Dimidium-ii. n. 2=a
half (subst).

Drachma-æ. f. 1=a drachm

Drachmæ duæ, tres, etc.=two, three, etc.,
drachms.

Semidrachma or drachma dimidia = half a
drachm (3ss).

Drachma cum semisse = a drachm and a half (3iiss).

Gramma-atis. n. 3=a gramme (gm.).

Granum-i. n. 2 = a grain (gr.).

Semigranum or granum dimidium=half a grain
(gr. ss.)

Grani pars quarta or quadrans=a quarter of a
grain (gr. $\frac{1}{4}$).

Grani pars tertia=a third of a grain (gr. $\frac{1}{3}$).

Grani pars quinta=a fifth of a grain (gr. $\frac{1}{5}$).

Grani partes quartas, tres, or q. tres quadrantes
=three-quarters of a grain (gr. $\frac{3}{4}$).

Grani partes tertias, duas=two-thirds of a grain (gr. $\frac{2}{3}$).

Grani partes quintas, quattuor=four-fifths of a grain (gr. $\frac{4}{5}$), etc.

Granum cum semisse=a grain and a half (gr. iss).

Grana duo, tria, etc.=two, three, etc., grains.

Gutta-æ. f. 1=a drop.

Libra-æ. f. 1=a pound (lb.)

Libræ dimidium or libra dimidia=half a pound.

Metrum-i. n. 2=a metre.

Centimetrum= $\frac{1}{100}$ of a metre. Cubic centimetres (cc).

Minimum-i. n. 2=a minim (℥.), *pl* minima.

Octarius-ii. m. 2=a pint (20 ounces), (Oj).

Quadrans-antis m. 3=a quarter (subst.)

Scrupulus-i. m. 2=a scruple (20 grains).

Semi. = half (adj.), semis-issis m. 3. = a half (subs.).

Uncia-æ. f. 1=an ounce ($\frac{1}{2}$).

Semiuncia, semuncia, uncia dimidia or uncia dimidium=half an ounce ($\frac{1}{4}$ ss).

Uncia cum semisse or sesuncia=an ounce and a half ($\frac{1}{2}$ iss)

Unciæ duæ, tres, etc.=two, three etc., ounces.

Example of a prescription fully written out:—

R Sodii Bicarbonatis drachmas duas ($\frac{1}{2}$ ii).

Magnesi Carbonatis drachmam ($\frac{1}{2}$ i).

Levis

Bismuthi Carbonatis grana viginti (gr. xx).

Acidi Hydrocyanici minima decem (℥. x).

diluti

Syrupi unciam cum semisse ($\frac{1}{2}$ iss)

Aquæ Chloroformi ad uncias sex ($\frac{1}{2}$ vi).

Fiat mistura. Solve et misce.

Signetur (or signa)—Capiat cochleare magnum ter in die ex aque cyatho parvo post cibos.

(Take a tablespoonful thrice a day in a wineglass of water after meals).

VOCABULARY OF LATIN WORDS AND PHRASES COMMONLY USED.

Note.

Nouns are given with their gender indicated by the letters m, f., n., and the declensions by the figures 1, 2, 3, 4, and 5.

Adjectives belong to the first three declensions only, and agree with their substantives in number, gender, and case.

Prepositions have the case which they govern indicated thus—(acc.), (gen.), (abl.).

Verbs have only the parts commonly employed given, with the translation as used in medicine. Verbs in the active voice put the object in the accusative case; and in the passive voice are related to the nominative. The conjugations are indicated by the figures 1, 2, 3, 4; deponent verbs have a passive form and active meaning.

The parts commonly used are:—

1. 3rd p. sing. or pl., pres. conj. act., e.g. *capiat-ant*—let (him or them) take, or take.
2. 3rd p. sing. or pl., pres. conj. pass, e.g. *capitur or antur*—let (it or them) be taken.
3. 3rd p. sing. or pl., fut. perf., ind. act., e.g. *convalescrit or erunt*—(he or they), shall have recovered.
4. 3rd p. sing. or pl., perf. conj. pass, e.g. *solutus sit or soluti sint*—(it or they) may have been dissolved.
5. 2nd p. imper. act., e.g. *recipe*—take (thou)
6. Pres. particp. act., e.g. *dolens-entis*—paining or painful.

7. *Past particip. pass.*, e.g. *dilutus-a-um*=
diluted.

8. *Gerundive. pass.*, e.g. *capiendus-a-um*=
to be taken

The participles and the gerundive agree with the substantive in number, gender, and case, and may be used as adjectives.

*For numerals see page 133, and for weights
and measures see page 135.*

Ad=to (acc.), e.g. *aquæ ad*, water to.

Adde 3.=add (thou), e.g. *solve et adde*, dissolve
and add.

Eger. m. ægra. f.=sick (person), the patient

Equalis-e.=equal, e.g. *æquales partes*, equal parts.

Agitatus-a-um.=shaken, e.g. *phiala prius agitata*,
shake the bottle (P.P.A.)

Affectus-a-um.=affected, e.g. *affectis partibus*, to
the parts affected.

Alternus-a-um=alternate, e.g. *diebus alternis*, every
other day.

Alvus i. f.=the bowels or stomach, e.g. *alvo jejuna*,
on an empty stomach.

Amplus-a-um.=large, e.g. *cyathus amplus*, a
tumblerful.

Ante.=before (acc.), e.g. *ante cibum*, before food.

Antea=formerly.

Applicandus-a-um 1. =to be applied.

Applicet. 1.=let (him) apply. *applicetur* or *entur*.
let (it or them) be applied.

Applicatio onis. f. 3.=an application.

Aqua-æ f. 1. =water, e.g. *ex aquâ*, in water.

Astrictus-a-um.=confined, e.g. *alvo astricta*, if con-
stipated.

Augendus-a-um.=to be increased, e.g. *augenda
dosis*, the dose to be increased.

Auris-is. f. 3.=the ear, e.g. *aures pone*, behind the
ears.

Bene. =well, e.g. *misce bene*, mix well.

- Bulliens-tis. =boiling, *e.g.*, ex aquā bulliente, in boiling water.
 Calidus-a-um. = hot, *e.g.* ex aquā calidā, in hot water
 Capiat or ant. 3. =let (him or them) take, or take.
 Capiatur or antur. =let (it or them) be taken.
 Capiendus-a-um. =to be taken.
 Cataplasma-atis. n. 3 =a poultice.
 Charta-æ. f. 1. =a paper, *e.g.* mitte chartas sex, send six (papers or) powders
 Cibus-i. m. 2 =food (pl.) meals, *e.g.* post cibum, after food or post cibos, after meals
 Cœna-æ. f. 1. =supper, *e.g.* post cœnam, after supper
 Collunarium-i. n. 2. =a nasal douche.
 Collutorium-i. n. 2. =a mouth wash.
 Communis-is-e. =common., *e.g.* pro potu communi, as a common drink.
 Collyrium-i. n. 2. =an eye wash.
 Cor-dis. n. 3. =the heart.
 Cras. = to-morrow, *e.g.* cras mane, to-morrow morning.
 Cum. =with (abl.), *e.g.* granum cum semisse, a grain and a half.
 Cutis-is. f. 3. =the skin, *e.g.* cutis integra, the unbroken skin.
 Dandus-a-um. 1. =to be given.
 Deinde =then or afterwards.
 Detur or dentur. 1. =let (it or them) be given.
 Dies-ei. m. 5. =a day, *e.g.* bis in die, twice a day.
 Dictus-a-um. =said, *e.g.* ut dictum, as said or prescribed.
 Dilutus-a-um. =diluted.
 Divide =divide. 3., *e.g.* divide in duas pilulas, divide into two pills.
 Dolens-entis. =painful, *e.g.* parti dolenti, to the painful part.
 Dolor-oris. f. 3. =pain, *e.g.* dolore urgenti. if the pain is severe.
 Donec =until, *e.g.* donec alvus soluta sit, until the bowels have acted.

Dorsum-i. n. 2.=the back, *e.g.* supra dorsum, over the back.

Dosis-is. f. 3.=(acc. dosin, abl. dosi) a dose.

Dum.=whilst.

Electuarium-i. n. 2.=an electuary.

Emplastrum-i. n. 2.=a plaster, *e.g.* emplastrum lyttæ, blistering plaster.

Enema-atis. n. 3.=a clyster or injection.

Et.—and., *e.g.* solve et misce, dissolve and mix

Ex.=out of or in (abl.), *e.g.* ex aquâ, in water.

Fauces-um. (pl.) f. 3.=the throat or tonsils.

Febris-is. f. 3.=fever, *e.g.* absente febris, if without fever.

Fervens-entis.=hot, *e.g.* ex aquâ fervente, in hot water.

Fiat or Fiant (Dep).=let (it or them) be made, make.

Flatus-ûs. m. 4.=flatulence.

Fluidus-a-um.=fluid.

Fortiter.=strongly.

Frustillum-i. m. 2.=a small piece, *e.g.* frustillo sacchari, on a lump of sugar.

Gargarisma-atis. n. 3.=a gargle.

Guttur-uris. n. 3. the throat.

Habeat=let (him) take, or give.

Habeatur or antur. 2. let (it or them) be taken, or given.

Habendus-a-um. =to be taken or given.

Haustus-i. m. 2.—a draught.

Hora-æ. f. 1.—an hour, *e.g.* horâ somni, the hour of sleep, bedtime.

In.=in (abl.) or into (acc.), *e.g.* in oculos, into the eyes.

Infra. =below (acc.), also adverb, beneath.

Infricandus-a-um. 1.=to be rubbed in.

Infricetur or entur. 1.=let (it or them) be rubbed in.

Infusum-i. (pl. infusa). n. 2.=an infusion.

Inhalatio-onis. f. 3.—an inhalation.

Inhalet. 1.=let (him) inhale.

Inhaletur or entur.=let (it or them) be inhaled.

Injectio-onis. f. 3.=an injection.

Instillandus-a-um. 1.=to be dropped in.

Instilletur or entur. 1.=let (it or them) be dropped in.

Integer-ra-rum.=whole, unbroken (see Cutis).

Inter.=between, among (acc.), *e.g.* inter cibos, between meals.

Intra.=inside, within (acc.), *e.g.* intra orem, inside the mouth.

Jejunus-a-um.=fasting, *e.g.* alvo jejuna, on an empty stomach.

Jentaculum-i. n. 2.=breakfast.

Lac-tis. n. 3.=milk, *e.g.* ex lacte, in milk.

Largus-a-um.=large, *e.g.* tria cochlearia larga, three tablespoonfuls.

Leniatur. 4.=may be relieved, *e.g.* donec dolor leniatur, until the pain is relieved.

Leniter=gently, *e.g.* tere leniter, rub gently.

Levatus-a-um.=relieved, *e.g.* donec levatus sit, until relieved.

Libet.=it pleases *e.g.* quantum libet, as much as you like (q.l.)

Linimentum-i. n. 2.=a liniment.

Linteum-ei. n. 2.=linen, *e.g.* linteum carptum, lint.

Liquidus-a-um.=liquid or loose, *e.g.* sedes liquidas, loose motions.

Liquor-oris. m. 3.=a fluid or liquor.

Lotio-onis. f. 3.=a lotion.

Lux-ucis. f. 3.=light, *e.g.* lucis prima, early in the morning.

Lytta-æ. f. 1.=a blister.

Magnus-a-um.=great, *e.g.* cochleare magnum, a tablespoonful.

Mane morning, *e.g.* primo mane, the first thing in the morning.

Massa-æ. f. 1.=a mass.

Medius-a-um.=medium, *e.g.* coch. medium, a dessertspoonful.

Meridies-ei m. 5.=noon, *e.g.* ante meridiem, a.m., post meridiem, p.m.

- Minimus-a-um. = smallest, *e.g.* coch. minimum, a teaspoonful.
 Minor-oris. = less, *e.g.* capiat dosin minorem, take a smaller dose.
 Misce. = mix (thou).
 Mistura-æ. f. 1. = a mixture.
 Mitigatus-a-um. = relieved, *e.g.* donec mitigatus sit, until relieved.
 Mitte. = send (thou).
 Mos-moris. m. 3. = manner, *e.g.*, more dicto, as directed.
 Nausea-æ. f. 1. = sickness, *e.g.* nauseâ urgente, if the sickness is troublesome.
 Nebula-æ. f. 1. = a spray.
 Necessitas-atis. f. 3. = necessity, *e.g.* si necessitas sit, if required.
 Nisi. = unless, *e.g.* nisi opus sit, unless required.
 Nox-tis. f. 3. = night, *e.g.* nocte maneque, night and morning.
 Oculus-i. m. 2 = the eye.
 Omnis-is-e. = all or every, *e.g.* omnibus horis, every hour.
 Ope. = by means of, *e.g.* ope pencilli, with a small brush.
 Optime. = well, *e.g.* tere optime simul, rub well together.
 Opus. = necessity, *e.g.* si opus sit, if necessary.
 Os-oris. n. 3. = the mouth, per orem, by the mouth.
 Pars-tis. f. 3. = a part, *e.g.* parti effectæ, to the part affected.
 Parvus-a-um. = small, *e.g.* coch. parvum, a teaspoonful.
 Pasta-æ. f. 1. = a paste.
 Paucus-a-um. = little or few, *e.g.* pauca: guttæ, a few drops.
 Paullum or Pauxillum-i. n. 2. = a little.
 Pectus-oris. n. 3. = the chest.
 Penicillum-i. n. 2 = camel's hair brush.
 Per. through, during (acc.), *e.g.* per anum, by the anus.

Persistat. = let (him) continue, *e.g.* persistat in usu
mistræ, continue the mixture.

Pessus-i. m. 2. = a pessary.

Phiala-æ. f. 1. = a bottle, *e.g.* phiala prius agitata,
shake the bottle (P.P.A.)

Pigmentum-i. n. 2. = paint.

Pilula-æ. f. 1. = a pill.

Pinge. 3. = paint (thou.)

Plene. = freely, *e.g.* plene dilutus, freely diluted.

Plenus-a-um. = full or large, *e.g.* coch. plenum, a
tablespoonful.

Pone. = behind (acc.), *e.g.* pone aures, behind the
ears.

Pone. 3. = place (thou.)

Post. = after (acc.), *e.g.* post prandium, after dinner.

Postea = afterwards.

Potus-ūs. m. 4. = a drink, *e.g.* pro potu communi,
for an ordinary drink.

Prandium-ii. n. 2. = dinner.

Primus-a-um. = first, *e.g.* primo mane, the first
thing in the morning.

Prius. = previously (see under Phiala.)

Pro. = before, for, or on behalf of (abl.), *e.g.* pro re
nata, occasionally.

Pulvis-eris. m. 3. = a powder.

Quando, = when, quodocunque, whenever.

Quantum = as much as, *e.g.* quantum sufficiat,
as much as required (q.s.)

Que. = and (enclitic), *e.g.* nocte maneque, night and
morning.

Qui-quæ-quod. = who or which. Cui, to which.
Cujus, of which.

Quisque, quaque, quodque. = each, every, *e.g.*
quartâquâque horâ, every four hours.

Quotidie. = daily.

Recentis-entis. = recent, fresh, *e.g.* infusum recens,
a fresh infusion.

Recipe. 3. = take (thou.)

Recuperatus-a-um. = recovered, *e.g.* quandoeger
recuperatus sit, when the patient has recovered.

Regio-ionis. f. 3. = the region, *e.g.* regioni cordis, over the heart.

Repetendus-a-um. 3. = to be repeated.

Repetat-ant, repetatur or antur. 3. = let (him or them) repeat, or let (it or them) be repeated.

Res-rei. f. 5. = a thing (see under Pro.)

Residuum-ui. n. 2. = the remainder.

Responderit or erint. 2. = (it or they) shall have acted.

Sæpe = often *e.g.* utendus sæpe, use often.

Saccharum-i. m. 2. = sugar (see frustillum.)

Satis. = enough.

Secundum. (acc.) = according to, *e.g.* secundum artem, with skill.

Sed. = but.

Sedes-is. f. 3. = a motion, *e.g.* post liquidas sedes, after loose motions.

Semper. = always.

Sensim. = gradually.

Septimana-æ. f. 1. = a week.

Sequens-entis. = following, *e.g.* die sequente, on the following day.

Si. = if, *e.g.* si opus sit, if necessary.

Signa or Signetur. 1. = label, or let (it) be directed.

Simul. = together. *e.g.* tere simul, rub together.

Sine. = without (abl.), *e.g.* sine cibo, without food, or fasting.

Singulus-a-um. = each, *e.g.* singula quaque dosi, with each dose.

Sit or sint. (Aux.). = (it or they) may be.

Solitus-a-um. = usual, *e.g.* more solito, in the usual manner, as usual.

Solutio-onis. f. 3. = a solution.

Solutus-a-um. = dissolved, *e.g.* quando solutus sit, when dissolved.

Solve. 3. = dissolve (thou).

Somnus-i. m. 2. = sleep, *e.g.* horâ somni, at bedtime.

Spongiola-æ. f. 1. = a small piece of sponge.

Statim. = immediately, *e.g.* utendus statim, use at once.

- Sudor-oris. m. 3. = sweating or perspiration.
- Sufficiat. 3. = (it) may suffice, *e.g.* quantum sufficiat as much as may suffice.
- Sumat or ant, sumatur or antur. 3. = let (him or them) take, or let (it or them) be taken.
- Sumendus-a-um. 3. = to be taken.
- Super. = on or over (acc.), *e.g.* super linteam carptum, on lint.
- Supervenerit or erint - (it or they) should have occurred.
- Supra. = over, above (acc.), *e.g.* ut supra, as above.
- Syrupus-i. m. 2. = syrup.
- Talis-is-e. = such, *e.g.* mitte tales xii., send twelve such. (Talia, n pl.).
- Tantum-i. n. 2. = as much, *e.g.* cum tanto aquæ, with as much water.
- Tepidus-a-um. = lukewarm, *e.g.* tepida aqua, tepid water.
- Tere. 3. = grind or rub (thou), *e.g.* tere simul optime, rub well together.
- Tunicatus-a-um. = coated or varnished, *e.g.* pilulæ tunicate, varnished pills.
- Tussis-is. f. 3. = a cough, *e.g.* tussi dolente, if the cough is painful.
- Ubi. = where.
- Unguentum-i. n. 2. = an ointment.
- Urgens-tis. = severe, *e.g.* urgenti dolore, if the pain is severe.
- Usus-ûs. m. 4. = use, *e.g.* pro usû externo, for outward use.
- Ut. = as or so as, *e.g.* ut dictum, ut fiat, as directed, so as to make.
- Utat or ant, utatur or antur. (dep.) = let (him or them) use, or let (it or them) be used.
- Utere. 3. (gov. abl.) = use thou, *e.g.* utere lotione, use the lotion.
- Utendus-a-um. = to be used, *e.g.* utendus statim to be used at once.
- Vapor-oris. f. 3. = vapour, or an inhalation.
- Ve (enclitic) or *e.g.* bis terve twice or thrice.

Vel. = or, *e.g.* nocte vel mane, night or morning.

Volueris. (irreg.) = you may wish, *e.g.* quantum volueris (q.v.), as much as you like.

Vomitus-ûs. m. 4. = vomiting.

Vulnus-eris (pl. vulnera), n. 3. = a wound.

ABBREVIATIONS COMMONLY USED IN PRESCRIPTIONS.

a. = ante, before.

a.c. = ante cibum or cibos, before food or meals.

ââ. = ana, of each.

ad lib. = ad libitum at pleasure.

æ.p. = æquales partes, equal parts.

aq. = aqua, water (ex. aq., in water)

b. = bis, twice.

b.i.d., b.d. or b.d.d. = bis in die, bis die, or bis de die, twice a day.

c. = cibum or cibos, food or meals.

c̄. = cum, with.

c̄. sem. = cum semisse, and a half.

cap. = capiat, take, or capiatur, or capiendus.

coch. = cochleare, spoonful.

coch. mag., med., or min. = a table, dessert, or teaspoonful.

d. = dies, a day.

D. or Div. = divide, divide.

F. or Ft. = fiat, make, or fiant.

Fl. = fluidus, fluid.

gtt. = guttae, drops.

h. or hab. = habeat, take, or habeatur or habendus.

h. = hora, an hour.

h.d. = horâ decubitus, at bedtime.

h.s. = horâ somni, at bedtime.

H.S.S. = horâ somni sumendus, to be taken at bedtime.

i. = in, in (abl.), or into (acc.), *e.g.* b.i.d.

l. = lac-lactis, milk (ex. l. = in milk.)

liq. = liquor, liquor.

- M. = misce, mix (thou).
 m. or mist. = mistura, a mixture.
 m. = mane, morning.
 m. p. = mane primo, early in the morning.
 m. d. = more dicto, as directed.
 Mt. or Mitt. = Mitte, send.
 n. = nocte, night (n. et m., night and morning).
 o. or om. = omni, every.
 o. h., o. n., or o. m. = omni horā, nocte, or mane,
 every hour, night, or morning.
 p. or pil. = pilula, a pill.
 p. or pulv. = pulvis, powder.
 p. c. = post cibum or cibos, after food or meals.
 p. prand = post prandium, after dinner.
 P.P.A. = phiala prius agitata, shake the bottle.
 p. r. n. = pro re nata, occasionally.
 q.l. = quantum libet, enough.
 q.s. = quantum sufficiat, sufficient.
 q.v. = quantum volueris, as much as you wish.
 rep. = repetat, repeat, or repetatur, or repetendus.
 s. or sum. = sumat, take, or sumatur or sumendus.
 S. = solve, dissolve (s. et m., dissolve and mix).
 s. a. = secundam artem, according to practice,
 with skill.
 ss. = semisse, half.
 s. or st. = statim, immediately.
 S. or Sig. = signa or signetur, label.
 s. o. s. = si opus sit, if necessary.
 t. = tales, of such.
 t. = ter, thrice.
 t. i. d., t. d. or t.d.d. = ter in die, ter die, or ter de
 die, thrice a day.
 t. d. s. = ter die sumendus, take thrice a day.
 Tr. = tinctura, a tincture.
 u. = ut, as or so as.
 u. d. = ut dictum, as directed.
 u. ft. = ut fiat, so as to make.
 2tis = secundis horis, every two hours.
 3tis = tertiis horis, every three hours.
 4tis = quartis horis, every four hours.
 6tis = sextis horis, every six hours.

INDEX.

A

- Abscess, 312.
 Acidity, 1 to 4, 39, 42 to 44, 457.
 Acne Rosacea, 316 to 318.
 ,, Vulgaris, 313 to 315.
 After Pains, 402, 403.
 Alcoholism (acute), 197 to 200.
 ,, (chronic), 201, 202.
 Alopecia, 319 to 324.
 Amenorrhœa, 404 to 407.
 Anæmia, 80 to 82, 213, 214.
 ,, (pernicious), 83, 84.
 Aneurysm, 88, 89.
 Angina Pectoris, 90 to 94.
 Aphæ or Stomatitis, 5 to 7, 189, 190, 42.
 Arsenical Poisoning, 203.
 Asthma, 109, 245 to 251, 254.

B

- Bladder (irritable), 381 to 383.
 Boils, 325, 326.
 Bronchial Asthma, 252 to 254.
 Bronchiectasis, 269, 270.
 Bronchitis (acute), 255 to 257, 430 to 432.
 ,, (chronic), 258 to 268, 433, 434.
 Burns, 327, 328.

C

- Cancer, 329, 330, 408, 409.
 Calculus (urinary), 384, 385.
 Catarrh, 271 to 273, 435 to 437.
 Cervicitis, 410.
 Chlorosis, 85 (see Anæmia).
 Cholera, 148 to 151.

- Chorea, 207, 208, 438, 439.
Colic (intestinal), 8, 9.
,, (lead), 25, 26, 204.
Constipation (acute), 23 to 26, 444, 445.
,, (chronic), 10 to 22, 35, 63 to 65, 441
to 443.
Convulsions (infantile), 440.
Corns, 331.
Cough, 274 to 279, 286, 287.
Croup, 446 to 448.
Cystitis, 386 to 390.

D

- Debility, 121, 209, 210, 259, 311, 449
Delirium, 197.
Dentition, 450.
Diarrhœa, 27 to 33, 195, 196, 293, 294, 451 to 456.
Diphtheria, 152 to 154.
Dropsy (cardiac), 95 to 98.
,, (renal), 391 to 393.
Dysentery, 155 to 158.
Dysmenorrhœa, 411 to 413.
Dyspepsia, 31, 34 to 44, 457, 458.
Dysphagia, 303.

E

- Eurache, 128, 129.
Eczema, 332 to 341.
Endocarditis, 99, 100.
Enteritis (infantile), 454.
Epilepsy, 211 to 214.
Erysipelas, 159 to 162.
Eye (inflammation), 130 to 135

F

- Favus, 459.
Fever (pyrexia), 175 to 181, 306 to 308, 460 to 463
Flatulence, 34, 45 to 49, 455, 456, 464, 465.
Fœtid Feet, 342

G

- Gall Stones, 50.
 Gastralgia, 51, 52.
 Gastric Ulcer, 53, 54.
 Gonorrhœa, 163 to 166.
 Gout (acute), 107 to 111.
 „ (chronic), 112 to 116, 309.

H

- Hæmatemesis, 55 to 61.
 Hæmaturia, 394.
 Hæmophilia, 86.
 Hæmoptysis, 280, 295, 296.
 Hæmorrhage (typhoid), 193, 194.
 Hæmorrhoids, 62 to 68.
 Hay Fever, 136 to 138, 167.
 Heart Disease, 101 to 105.
 Hepatic Dyspepsia, 36, 40, 450.
 Herpes Zoster, 343 to 345.
 Hiccough, 215.
 Hysteria, 216 to 218, 233, 407.

I

- Incontinence of Urine, 395 to 397, 467.
 Influenza, 168.
 Insomnia, 199, 203, 219 to 225, 468 to 470.
 Intermittent Fever, 160.

L

- Laryngitis, 281 to 284, 300 to 302.
 Lead Poisoning, 25, 26, 204.
 Leprosy, 170.
 Leucorrhœa, 414, 415.
 Lumbago, 126, 127.
 Lupus, 346 to 349.

M

- Malaria, 171 to 172.
 Marasmus, 466.

- Menorrhagia, 416.
Migraine, 229, 230.
Morphia or Opium Poisoning, 205.

N

- Nephritis, 398 to 401.
Nervous Excitability, 226
Neuralgia, 227 to 237.
Neurasthenia, 238 to 240.
Neuritis, 241, 242.
Night Sweats, 291, 292.
Nipples (sore), 417.
Nose and Throat Inflammations, 136 to 138, 167, 273.

O

- Œdema of the Lung, 285, 471.
Opium or Morphia Poisoning, 205.
Orchitis, 418 to 420.
Ovaritis or Pelvic Cellulitis, 421.

P

- Paralysis, 243, 244.
Pediculosis, 350, 351.
Pelvic Cellulitis, 421
Pericarditis, 106.
Phthisis, 286 to 303, 473.
Plague, 174.
Pleurisy, 304, 305.
Pneumonia, 306 to 311, 472.
Pruritis, 355 to 358, 422 to 425.
Psoriasis, 352 to 354.
Purpura, 87, 359.
Pyrexia (fever), 175 to 181, 460 to 463.

R

- Relapsing Fever, 182.
Rheumatism (acute), 117 to 119.
" (chronic), 120 to 127.
Rigidity of Os uteri, 426.
Ringworm, 360 to 363, 459.
Round Worms, 483, 484.

S

- Sea Sickness, 71.
 Scabies, 364 to 368.
 Scarlet Fever, 183, 184.
 Sickness, 69, 70, 198.
 „ (to produce), 77 to 79, 446, 447.
 Sleeplessness (insomnia), 199, 208, 219 to 225. 468
 to 470.
 Strychnine Poisoning, 206.
 Syphilis (constitutional), 185 to 190.
 Syphilitic Eruptions, 369 to 372.
 Subinvolution, 427.

T

- Tabes Mesenterica, 474, 475.
 Tape Worms, 72 to 74, 487.
 Tetanus, 191, 192.
 Thread Worms, 75, 483, 485, 486.
 Throat Inflammations, 136 to 138.
 „ (relaxed), 139 to 142.
 Tonsillitis, 143 to 147.
 Tympanitis, 76.
 Typhoid Fever, 193 to 196.

U

- Ulcers, 376 to 378.
 Urticaria, 379, 380.

V

- Vaginismus, 428.
 Vomiting (to produce), 77 to 79, 446, 447.

- Warts, 373, 374.
 Whooping Cough, 476 to 482.
 Worms, 72 to 75, 483 to 487.
 Wounds (superficial), 373.



4/14

Sydney 7/1/2000

2-10-2000

